

#HW017

Histone H4R3me2a Polyclonal Antibody

Catalog: HW017

100µl

Orders: order@signalwayantibody.com

Storage: Store at -20°C or -80°C.

Support: tech@signalwayantibody.com

Avoid freeze / thaw cycles.

Web: www.sabbiotech.com

Application	Species Reactivity	Source	Molecular Wt.
WB IF IP ChIP	Hu Mu Rt Other(Wide range)	Rabbit Polyclonal Ab	15kDa

Description: Antibodies were produced by immunizing rabbits and were purified by antigen affinity-chromatography.

Immunogen: A synthetic methylated peptide corresponding to residues surrounding Arg3 of human histone H4

Formulation: Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

Synonyms: H3t; H3.4; H3/g; H3FT;

Accession No.: Gene ID: 8290 Swiss Prot: Q16695

Background:

Histones are basic nuclear proteins that are responsible for the nucleosome structure of the chromosomal fiber in eukaryotes. Nucleosomes consist of approximately 146 bp of DNA wrapped around a histone octamer composed of pairs of each of the four core histones (H2A, H2B, H3, and H4). The chromatin fiber is further compacted through the interaction of a linker histone, H1, with the DNA between the nucleosomes to form higher order chromatin structures. This gene is intronless and encodes a member of the histone H4 family. Transcripts from this gene lack polyA tails; instead, they contain a palindromic termination element.

Application:


Recommended Dilutions:

WB 1:500 - 1:2000


IF 1:50- 1:200

IP 1:50- 1:100


ChIP 1:50- 1:200



Western blot analysis of extracts of HeLa cell line and H3 protein expressed in E.coli., using H4R3 me2a antibody.



Dot-blot analysis of all sorts of methylation peptides using H4R3 me2a antibody.



Immunofluorescence analysis of 293T cell using H4R3me2a antibody. Blue: DAPI for nuclear staining.

This product is for in vitro research use only and is not intended for use in humans or animals.