PGA5 Antibody

Catalog No: #37105



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| $\overline{}$ | | | |
|------------------|------|-----|------|
| | escr | מוי | tion |
| \boldsymbol{L} | COUL | ıv | เเบเ |

| Product Name | PGA5 Antibody |
|-----------------------|---|
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antigen affinity purification. |
| Applications | WB IHC |
| Species Reactivity | Hu |
| Specificity | The antibody detects endogenous levels of total PGA5 protein. |
| Immunogen Type | Peptide |
| Immunogen Description | Synthetic peptide corresponding to a region derived from internal residues of human Pepsinogen 5, group I |
| | (pepsinogen A) |
| Target Name | PGA5 |
| Other Names | PGA3; PGA4 |
| Accession No. | Swiss-Prot#: P0DJD9NCBI Gene ID: 5222Gene Accssion: NP_055039 |
| Uniprot | P0DJD9 |
| GeneID | 5222; |
| SDS-PAGE MW | 42kd |
| Concentration | 2.1mg/ml |
| Formulation | Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol. |
| Storage | Store at -20°C |
| | |

Application Details

Western blotting: 1:500-1:2000
Immunohistochemistry: 1:25-1:100

Images




Gel: 10%SDS-PAGE

Lysates (from left to right): Human liver cancer tissue

Amount of lysate: 60ug per lane Primary antibody: 1/1000 dilution Secondary antibody dilution: 1/8000

Exposure time: 10 minutes



Immunohistochemical analysis of paraffin-embedded Human gastric cancer tissue using #37105 at dilution 1/60.

Background

The genes coding for human pepsinogen (PGA3, PGA4, and PGA5) were assigned to chromosome region 11q13 by in situ hybridization. Previously we localized the PGA gene complex to a centromeric region of chromosome 11 (p11----q13) by Southern blot analysis of mouse-human somatic cell hybrids.? Shows particularly broad specificity; although bonds involving phenylalanine and leucine are preferred, many others are also cleaved to some extent.

Note: This product is for in vitro research use only