

## DCAF6 Antibody

Catalog No: #37778

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)


## Description

|                       |  |
|-----------------------|--|
| Product Name          | DCAF6 Antibody   |
| Host Species          | Rabbit   |
| Clonality             | Polyclonal   |
| Purification          | Antigen affinity purification.   |
| Applications          | IHC  |
| Species Reactivity    | Hu Ms  |
| Specificity           | The antibody detects endogenous levels of total DCAF6 protein.   |
| Immunogen Type        | Peptide  |
| Immunogen Description | Synthetic peptide corresponding to residues near the C terminal of human DDB1 and CUL4 associated factor 6 |
| Target Name           | DCAF6  |
| Other Names           | NRIP; ARCAP; IQWD1; PC326; MSTP055; 1200006M05Rik  |
| Accession No.         | Swiss-Prot#: Q58WW2NCBI Gene ID: 55827Gene Accssion: NP_001017977  |
| Uniprot               | Q58WW2   |
| GeneID                | 55827;   |
| Concentration         | 1.7mg/ml   |
| Formulation           | Rabbit IgG in pH7.4 PBS, 0.05% NaN <sub>3</sub> , 40% Glycerol.  |
| Storage               | Store at -20°C   |


## Application Details

Immunohistochemistry: 1:50-1:200

## Images



Immunohistochemical analysis of paraffin-embedded Human thyroid cancer tissue using #37778 at dilution 1/30.



Immunohistochemical analysis of paraffin-embedded Human brain tissue using #37778 at dilution 1/30.

## Background

NRIP (Nuclear receptor interaction protein), also known as IQWD1 (IQ motif and WD repeat-containing protein 1), MSTP055, ARCAP or PC326, is an 860 amino acid protein that localizes to the nucleus and contains one IQ domain and seven WD-repeats. Expressed in testis, skeletal muscle, prostate and heart, NRIP functions as a ligand-dependent coactivator of nuclear receptors and specifically enhances the transcriptional activity of AR (androgen receptor) and GR (glucocorticoid receptor). NRIP exists as three isoforms that are produced by alternative splicing events.

Note: This product is for in vitro research use only