## WAVE1 (Phospho-Tyr125) Antibody

Catalog No: #12156

Package Size: #12156-1 50ul #12156-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

WAVE1 (Phospho-Tyr125) Antibody
Rabbit
Polyclonal
Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates.
Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho
specific antibodies were removed by chromatogramphy using non-phosphopeptide.
WB IHC
Hu Ms
The antibody detects endogenous levels of WAVE1 only when phosphorylated at tyrosine 125.
peptide
Peptide sequence around phosphorylation site of tyrosine 125 (E-T-Y(p)-D-V) derived from Human WAVE1.
WAVE1
Phospho
KIAA0269; SCAR1; Verprolin homology domain-containing protein 1; WAS1; WASF1; WASP-family protein
member 1; Wiskott-Aldrich syndrome protein family member 1
Swiss-Prot#:Q92558;NCBI Gene#:8936
Q92558
8936;
70kd
1.0mg/ml
Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% sodium azide
and 50% glycerol.
Store at -20°C

Application Details
Western blotting: 1:500~1:3
Immunohistochemistry: 1:50-

## Images



Western blot analysis of extracts from NIH/3T3 cells, treated with Insulin (0.01U/ml, 15mins), using WAVE1 (Phospho-Tyr125) antibody #12156. The lane on the right is treated with the synthesized peptide.



Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue using WAVE1 (Phospho-Tyr125) antibody #12156. The picture on the right is treated with the synthesized peptide.

## Background

Downstream effector molecule involved in the transmission of signals from tyrosine kinase receptors and small GTPases to the actin cytoskeleton. Promotes formation of actin filaments. Part of the WAVE complex that regulates lamellipodia formation. The WAVE complex regulates actin filament reorganization via its interaction with the Arp2/3 complex.

Note: This product is for in vitro research use only