Protocadherin-1 antibody

Catalog No: #22346



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| Description | Support: tech@signalwayantibody.com |
|-----------------------|--|
| Product Name | Protocadherin-1 antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Purified by antigen-affinity chromatography. |
| Applications | WB IHC |
| Species Reactivity | Hu |
| Immunogen Type | Recombinant protein |
| Immunogen Description | Recombinant protein fragment contain a sequence corresponding to a region within amino acids 710 and 941 |
| | of Protocadherin-1 |
| Target Name | Protocadherin-1 |
| Accession No. | Swiss-Prot:Q08174Gene ID:5097 |
| Uniprot | Q08174 |
| GeneID | 5097; |
| Concentration | 0.7mg/ml |
| Formulation | Supplied in 0.1M Tris-buffered saline with 20% Glycerol (pH7.0). 0.01% Thimerosal was added as a |
| | preservative. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |

Application Details

Predicted MW: 115kd


Immunohistochemistry: 1:100-1:250

Images



Sample (30 ug of whole cell lysate) A: HCT116 7.5% SDS PAGE

Primary antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded BT474 xenograft, using PCDH1 antibody at 1: 100 dilution.

Background

This gene belongs to the protocadherin subfamily within the cadherin superfamily. The encoded protein is a membrane protein found at cell-cell boundaries. It is involved in neural cell adhesion, suggesting a possible role in neuronal development. The protein includes an extracelllular region, containing 7 cadherin-like domains, a transmembrane region and a C-terminal cytoplasmic region. Cells expressing the protein showed cell aggregation activity. Alternative splicing occurs in this gene. [provided by RefSeq]

Note: This product is for in vitro research use only