p53(Phospho-Ser15) Antibody

Catalog No: #11094

Package Size: #11094-1 50ul #11094-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


| Description | | | | |
|-----------------------|--|--|--|--|
| Product Name | p53(Phospho-Ser15) Antibody | | | |
| Host Species | Rabbit | | | |
| Clonality | Polyclonal | | | |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. | | | |
| | Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho | | | |
| | specific antibodies were removed by chromatogramphy using non-phosphopeptide. | | | |
| Applications | WB IHC IF | | | |
| Species Reactivity | Hu | | | |
| Specificity | The antibody detects endogenous level of p53 only whenphosphorylated at serine15. | | | |
| Immunogen Type | Peptide-KLH | | | |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 15 (P-L-S(p)-Q-E) derived from Human p53. | | | |
| Target Name | p53 | | | |
| Modification | Phospho | | | |
| Other Names | Tumor suppressor p53; Phosphoprotein p53; Antigen NY-CO-13; TP53; | | | |
| Accession No. | Swiss-Prot: P04637NCBI Protein: NP_000537.3 | | | |
| Uniprot | P04637 | | | |
| GeneID | 7157; | | | |
| Concentration | 1.0mg/ml | | | |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.02% | | | |
| | sodium azide and 50% glycerol. | | | |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. | | | |
| | | | | |

| Application Details | | | |
|--------------------------------|----|--|--|
| Predicted MW: 53kd | | | |
| Western blotting: 1:500~1:1000 | | | |
| Immunohistochemistry: 1:50~1: | 00 | | |
| Immunofluorescence: 1:100~1:2 | 00 | | |


Images




Western blot analysis of extracts from 293 cells untreated or treated with UV using p53(Phospho-Ser15) Antibody #11094.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using p53 (Phospho-Ser15) Antibody #11094 (left) or the same antibody preincubated with blocking peptide #51094 (right).



Immunofluorescence staining of methanol-fixed Hela cells using p53(Phospho-Ser15) Antibody #11094.



Immunofluorescence staining of methanol-fixed Hela cells using p53 (Phospho-Ser15) Antibody #11094.

Background

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over.

Lin T, et al. (2005) Nat Cell Biol; 7(2): 165-71.

Vega FM, et al. (2004) Mol Cell Biol; 24(23): 10366-80.

Li J, et al. (2004) J Biol Chem; 279(40): 41275-9.

Wang J, et al. (2004) J Biol Chem; 279(38): 39584-92.

Note: This product is for in vitro research use only