

p53(Phospho-Thr18) Antibody

Catalog No: #11095

Package Size: #11095-1 50ul #11095-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	p53(Phospho-Thr18) Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide.
Applications	WB
Species Reactivity	Hu
Specificity	The antibody detects endogenous level of p53 only when phosphorylated at threonine 18.
Immunogen Type	Peptide-KLH
Immunogen Description	Peptide sequence around phosphorylation site of threonine 18 (Q-E-T(p)-F-S) derived from Human p53.
Target Name	p53
Modification	Phospho
Other Names	Antigen NY-CO-13; Phosphoprotein p53; TP53; Tumor suppressor p53;
Accession No.	Swiss-Prot: P04637NCBI Protein: NP_000537.3
Uniprot	P04637
GenelD	7157;
Concentration	1.0mg/ml
Formulation	Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 53kd

Western blotting: 1:500~1:1000

Images



Western blot analysis of extracts from HT29 and MDA cells untreated or treated with anisomycin using p53(Phospho-Thr18) Antibody #11095.

Background

Acts as a tumor suppressor in many tumor types; induces growth arrest or apoptosis depending on the physiological circumstances and cell type. Involved in cell cycle regulation as a trans-activator that acts to negatively regulate cell division by controlling a set of genes required for this process. One of the activated genes is an inhibitor of cyclin-dependent kinases. Apoptosis induction seems to be mediated either by stimulation of BAX and FAS antigen expression, or by repression of Bcl-2 expression. Implicated in Notch signaling cross-over.

Lin T, et al. (2005) *Nat Cell Biol*; 7(2): 165-71.

Vega FM, et al. (2004) *Mol Cell Biol*; 24(23): 10366-80.

Li J, et al. (2004) *J Biol Chem*; 279(40): 41275-9.

Wang J, et al. (2004) *J Biol Chem*; 279(38): 39584-92.

Note: This product is for in vitro research use only