

CD27L(CD27L(CD70)) antibody

Catalog No: #22409

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


| | |
|-----------------------|--|
| Product Name | CD27L(CD27L(CD70)) antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Purified by antigen-affinity chromatography. |
| Applications | WB |
| Species Reactivity | Hu |
| Immunogen Type | Peptide |
| Immunogen Description | Synthetic peptide contain a sequence corresponding to a region within amino acids 129 and 193 of CD27L (CD70) |
| Target Name | CD27L(CD27L(CD70)) |
| Accession No. | Swiss-Prot:P32970Gene ID:970 |
| Uniprot | P32970 |
| GeneID | 970; |
| Concentration | 1mg/ml |
| Formulation | Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a preservative. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |

Application Details

Predicted MW: 21kd

Western blotting: 1:500-1:3000

Images



Sample (30 ug of whole cell lysate)

A: JurKat

12% SDS PAGE

CD27L (CD70) antibody diluted at 1: 1000

Background

The protein encoded by this gene is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T and B lymphocytes. It induces proliferation of costimulated T cells,

enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis. [provided by RefSeq]

Note: This product is for in vitro research use only