## **EEPD1** Antibody

Catalog No: #47092



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_			
	escri	ınt	ion.
$\boldsymbol{L}$	COUL	Iυι	IUII

EEPD1 Antibody
Rabbit
Polyclonal
Antigen affinity purification
WB, IHC
Hu
The antibody detects endogenous levels of total EEPD1 protein.
peptide
Synthetic peptide of human EEPD1
EEPD1
HSPC107
Swiss-Prot#:Q7L9B9 NCBI Gene ID:80820Gene Accssion:NP_085139
Q7L9B9
80820;
63 kDa
0.7mg/ml
Rabbit IgG in pH7.4 PBS, 0.05% NaN3, 40% Glycerol.
Store at -20C


## Application Details

Western blotting:1:200-1000Immunofluorescence:1: 20-100

## **Images**



The image is immunohistochemistry of paraffin-embedded Human breast cancer tissue using 47092(EEPD1 Antibody) at dilution 1/20. (Original magnification: ?00)



Gel: 8%SDS-PAGE

Lysate: 40 µg, Lane 1-3: PC-3 cell, Human cerebrum tissue,

231 cell lysates

Primary antibody:EEPD1 Antibody at dilution 1/300 Secondary antibody: Goat anti rabbit IgG at 1/8000 dilution

Exposure time: 30 seconds

## Background

EEPD1(endonuclease/exonuclease/phosphatase family domain containing 1), also known as HSPC107, is a 569 amino acid protein that contains one HhH domain. A significant decrease in the relative transcriptional level of EEPD1 is induced by long-term heat stress exposure. Conversely, EEPD1 is up-regulated in bovine adipogenic processes related to intramuscular pre-adipocyte differentiation. Encoded by a gene that maps to human chromosome 7p14.2, EEPD1 plays a role in DNA binding and repair. Chromosome 7 makes up about 5% of the human genome and contains 158 million bases encoding more than 1,000 genes. Osteogenesis imperfecta, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome are associated with Chromosome 7.

Note: This product is for in vitro research use only