

Myosin Light Chain 2 (Phospho-Ser19) Antibody

Catalog No: #11114



Package Size: #11114-1 50ul #11114-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | Myosin Light Chain 2 (Phospho-Ser19) Antibody |
| Host Species | Rabbit |
| Clonality | Polyclonal |
| Purification | Antibodies were produced by immunizing rabbits with synthetic phosphopeptide and KLH conjugates. Antibodies were purified by affinity-chromatography using epitope-specific phosphopeptide. Non-phospho specific antibodies were removed by chromatography using non-phosphopeptide. |
| Applications | WB IHC |
| Species Reactivity | Hu Rt |
| Specificity | The antibody detects endogenous level of Myosin Light Chain 2 only when phosphorylated at serine 19. |
| Immunogen Type | Peptide-KLH |
| Immunogen Description | Peptide sequence around phosphorylation site of serine 19 (A-T-S(p)-N-V) derived from Human Myosin Light Chain 2. |
| Target Name | Myosin Light Chain 2 |
| Modification | Phospho |
| Other Names | LC20; MLC2; MRLC1; MYRL2; MLC-2C; MMIHS4 |
| Accession No. | Swiss-Prot: P24844 |
| Uniprot | P24844 |
| GeneID | 10398; |
| Concentration | 1.0mg/ml |
| Formulation | Supplied at 1.0mg/mL in phosphate buffered saline (without Mg ²⁺ and Ca ²⁺), pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. |
| Storage | Store at -20°C for long term preservation (recommended). Store at 4°C for short term use. |


Application Details

Predicted MW: 18kd


Western blotting: 1:500~1:1000

Immunohistochemistry: 1:50~1:100

Images



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using Myosin Light Chain 2 (Phospho-Ser19) Antibody #11114 (left) or the same antibody preincubated with blocking peptide (right).



Western blot analysis of extracts from 293 cells untreated or treated with starvation using Myosin Light Chain 2 (Phospho-Ser19) Antibody #11114.

Background

Myosin regulatory subunit that plays an important role in regulation of both smooth muscle and nonmuscle cell contractile activity via its phosphorylation. Implicated in cytokinesis, receptor capping, and cell locomotion

Janiak A, et al. (2006) *Mol Biol Cell*. Apr; 17(4): 1606-1619.

Croft DR, et al. (2006) *Mol Cell Biol*. 2 Jun; 26(12): 4612-4627

Li Z, et al. (2006) *Mol Cell Biol*. Jun; 26(11): 4240-4256

Note: This product is for in vitro research use only