

NAD-dependent deacetylase sirtuin-2

Catalog No: #AP78516



Package Size: #AP78516-1 50ug #AP78516-2 100ug #AP78516-3 1mg

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com

Description

Product Name	NAD-dependent deacetylase sirtuin-2
Brief Description	Recombinant Protein
Host Species	E.coli
Purification	Greater than 90% by SDS-PAGE
Species Reactivity	Human
Immunogen Description	65-340AA
Other Names	SIR2L, SIR2L2, SIR2-like protein 2
Accession No.	Q8IXJ6 Gene name: SIRT2
Uniprot	Q8IXJ6
GeneID	22933;
Calculated MW	30.36
Tag Info	His
Formulation	50mM NaH ₂ PO ₄ , 500mM NaCl Buffer with 500mM Imidazole, 10% glycerol (pH 8.0)
Storage	Store at -20C. (Avoid repeated freezing and thawing.) Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Background

NAD-dependent protein deacetylase, which deacetylates the 'Lys-40' of alpha-tubulin. Involved in the control of mitotic exit in the cell cycle, probably via its role in the regulation of cytoskeleton.

References

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Note: This product is for in vitro research use only