

## Catenin beta-1

Catalog No: #AP78747

Package Size: #AP78747-1 50ug #AP78747-2 100ug #AP78747-3 1mg

Orders: [order@signalwayantibody.com](mailto:order@signalwayantibody.com)Support: [tech@signalwayantibody.com](mailto:tech@signalwayantibody.com)

## Description

Product Name	Catenin beta-1
Brief Description	Recombinant Protein
Host Species	E.coli
Purification	Greater than 90% by SDS-PAGE
Species Reactivity	Human
Immunogen Description	654-781AA
Other Names	OK/SW-cl.35,PRO2286,CTNNB,Beta-catenin
Accession No.	P35222Gene name:CTNNB1
Uniprot	P35222
GeneID	1499;
Calculated MW	14.08
Tag Info	His
Formulation	50mM NaH <sub>2</sub> PO <sub>4</sub> , 500mM NaCl Buffer with 500mM Imidazole,10%glycerol(PH8.0)
Storage	Store at -20C. (Avoid repeated freezing and thawing.)Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

## Background

Key downstream component of the canonical Wnt signaling pathway. In the absence of Wnt, forms a complex with AXIN1, AXIN2, APC, CSNK1A1 and GSK3B that promotes phosphorylation on N-terminal Ser and Thr residues and ubiquitination of CTNNB1 via BTRC and its subsequent degradation by the proteasome. In the presence of Wnt ligand, CTNNB1 is not ubiquitinated and accumulates in the nucleus, where it acts as a coactivator for transcription factors of the TCF/LEF family, leading to activate Wnt responsive genes. Involved in the regulation of cell adhesion. Acts as a negative regulator of centrosome cohesion. Involved in the CDK2/PTPN6/CTNNB1/CEACAM1 pathway of insulin internalization. Blocks anoikis of malignant kidney and intestinal epithelial cells and promotes their anchorage-independent growth by down-regulating DAPK2. Disrupts PML function and PML-NB formation by inhibiting RANBP2-mediated sumoylation of PML (PubMed:17524503, PubMed:18077326, PubMed:18086858, PubMed:18957423, PubMed:21262353, PubMed:22647378, PubMed:22699938, PubMed:22155184). Promotes neurogenesis by maintaining sympathetic neuroblasts within the cell cycle.

## References

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Note: This product is for in vitro research use only