

Serine/threonine-protein kinase mTOR

Catalog No: #AP79038



Package Size: #AP79038-1 50ug #AP79038-2 100ug #AP79038-3 1mg

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Serine/threonine-protein kinase mTOR
Brief Description	Recombinant Protein
Host Species	E.coli
Purification	Greater than 90% by SDS-PAGE
Species Reactivity	Rat
Immunogen Description	Recombinant Protein
Other Names	Frap1,Raft1,FK506-binding protein 12-rapamycin complex-associated protein 1,FKBP12-rapamycin complex-associated protein,Mammalian target of rapamycin,Mechanistic target of rapamycin,Rapamycin target protein 1,mTOR,RAPT1
Accession No.	P42346Gene name:Mtor
Uniprot	P42346
GeneID	56718;
Tag Info	His
Formulation	50mM NaH ₂ PO ₄ , 500mM NaCl Buffer with 500mM Imidazole,10%glycerol(PH8.0)
Storage	Store at -20C. (Avoid repeated freezing and thawing.)Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Background

Serine/threonine protein kinase which is a central regulator of cellular metabolism, growth and survival in response to hormones, growth factors, nutrients, energy and stress signals. MTOR directly or indirectly regulates the phosphorylation of at least 800 proteins. Functions as part of 2 structurally and functionally distinct signaling complexes mTORC1 and mTORC2 (mTOR complex 1 and 2). Activated mTORC1 up-regulates protein synthesis by phosphorylating key regulators of mRNA translation and ribosome synthesis. This includes phosphorylation of EIF4EBP1 and release of its inhibition toward the elongation initiation factor 4E (eIF4E). Moreover, phosphorylates and activates RPS6KB1 and RPS6KB2 that promote protein synthesis by modulating the activity of their downstream targets including ribosomal protein S6, eukaryotic translation initiation factor EIF4B, and the inhibitor of translation initiation PDCD4. Stimulates the pyrimidine biosynthesis pathway, both by acute regulation through RPS6KB1-mediated phosphorylation of the biosynthetic enzyme CAD, and delayed regulation, through transcriptional enhancement of the pentose phosphate pathway which produces 5-phosphoribosyl-1-pyrophosphate (PRPP), an allosteric activator of CAD at a later step in synthesis, this function is dependent on the mTORC1 complex. Regulates ribosome synthesis by activating RNA polymerase III-dependent transcription through phosphorylation and inhibition of MAF1 an RNA polymerase III-repressor. In parallel to protein synthesis, also regulates lipid synthesis through SREBF1/SREBP1 and LPIN1. To maintain energy homeostasis mTORC1 may also regulate mitochondrial biogenesis through regulation of PPARGC1A. mTORC1 also negatively regulates autophagy through phosphorylation of ULK1. Under nutrient sufficiency, phosphorylates ULK1 at 'Ser-758', disrupting the interaction with AMPK and preventing activation of ULK1. Also prevents autophagy through phosphorylation of the autophagy inhibitor DAP. mTORC1 exerts a feedback control on upstream growth factor signaling that includes phosphorylation and activation of GRB10 a INSR-dependent signaling suppressor. Among other potential targets mTORC1 may phosphorylate CLIP1 and regulate microtubules. As part of the mTORC2 complex MTOR may regulate other cellular processes including survival and organization of the cytoskeleton. Plays a critical role in the phosphorylation at 'Ser-473' of AKT1, a pro-survival effector of phosphoinositide 3-kinase, facilitating its activation by PDK1. mTORC2 may regulate the actin cytoskeleton, through phosphorylation of PRKCA, PXN and activation of the Rho-type guanine nucleotide exchange factors RHOA and RAC1A or RAC1B. mTORC2 also regulates the phosphorylation of SGK1 at 'Ser-421'. Regulates osteoclastogenesis by adjusting the expression of CEBPB isoforms.

References

Repeated freezing and thawing is not recommended. Store working aliquots at 4°C for up to one week.

Note: This product is for in vitro research use only