

GPR73a Antibody

Catalog No: #47928

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	GPR73a Antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	The antibody was purified by immunogen affinity chromatography.
Applications	WB, IHC, IF/ICC
Species Reactivity	Hu,Ms,Rt
Specificity	Recognizes endogenous levels of GPR73a protein.
Immunogen Description	KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human GPR73a.
Target Name	PROKR1
Other Names	GPR73; PKR1; Prokineticin receptor 1; PK-R1; G-protein coupled receptor 73; G-protein coupled receptor ZAQ; GPR73a
Accession No.	Swiss-Prot#:Q8TCW9NCBI Gene ID:10887
Uniprot	Q8TCW9
GeneID	10887;
Calculated MW	45KD
Concentration	1 mg/ml
Formulation	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Storage	Store at -20°C


Application Details

WB (1/500 - 1/1000), IHC (1/50 - 1/100), IF/ICC (1/50 - 1/200)


Images



Immunofluorescent analysis of GPR73a staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).



Western blot analysis of GPR73a expression in BV2 (A), H9C2 (B), HCC827 (C), HEK293T (D) whole cell lysates



Immunohistochemical analysis of GPR73a staining in human liver cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Note: This product is for in vitro research use only