

GAPDH Antibody

Catalog No: #48102



Package Size: #48102-1 50ul #48102-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	GAPDH Antibody
Host Species	Mouse
Clonality	Monoclonal
Clone No.	1-G2
Purification	ProA affinity purified
Applications	WB, ICC
Species Reactivity	Hu, Ms, Rt, zebrafish
Immunogen Description	peptide
Other Names	38 kDa BFA-dependent ADP-ribosylation substrate antibody aging associated gene 9 protein antibody Aging-associated gene 9 protein antibody BARS-38 antibody cb609 antibody EC 1.2.1.12 antibody Epididymis secretory sperm binding protein Li 162eP antibody G3P_HUMAN antibody G3PD antibody G3PDH antibody GAPD antibody GAPDH antibody Glyceraldehyde 3 phosphate dehydrogenase antibody Glyceraldehyde-3-phosphate dehydrogenase antibody HEL-S-162eP antibody KNC-NDS6 antibody MGC102544 antibody MGC102546 antibody MGC103190 antibody MGC103191 antibody MGC105239 antibody MGC127711 antibody MGC88685 antibody OCAS, p38 component antibody OCT1 coactivator in S phase, 38-KD component antibody peptidyl cysteine S nitrosylase GAPDH antibody Peptidyl-cysteine S-nitrosylase GAPDH antibody wu:fb33a10 antibody
Accession No.	Swiss-Prot#:P04406
Uniprot	P04406
GeneID	2597;
Calculated MW	36 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details


WB: 1:2000-1:5,000 ICC: 1:200

Images




Western blot analysis of GAPDH on different cells lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room temperature.


Positive control:
 Lane 1: Hela cell lysate
 Lane 2: A549 cell lysate
 Lane 3: HepG2 cell lysate
 Lane 4: PC-12 cell lysate
 Lane 5: F9 cell lysate



ICC staining GAPDH in D3 cells (red). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS and counterstained with DAPI in order to highlight the nucleus (blue).



ICC staining of GAPDH in A431 cells (red). Formalin fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 10 minutes at room temperature and blocked with 1% Blocker BSA for 15 minutes at room temperature. Cells were probed with the primary antibody for 1 hour at room temperature, washed with PBS. Alexa Fluor 555 Goat anti-Mouse IgG was used as the secondary antibody at 1/100 dilution. The nuclear counter stain is DAPI (blue).



Western blot analysis of GAPDH on zebrafish tissue lysates. Proteins were transferred to a PVDF membrane and blocked with 5% BSA in PBS for 1 hour at room temperature. The primary antibody was used in 5% BSA at room temperature for 2 hours. Goat Anti-Mouse IgG - HRP Secondary Antibody at 1:5,000 dilution was used for 1 hour at room temperature.

Background

GAPDH (Glyceraldehyde-3-phosphate dehydrogenase) has both glyceraldehyde-3-phosphate dehydrogenase and nitrosylase activities, thereby playing a role in glycolysis and nuclear functions, respectively. It participates in nuclear events including transcription, RNA transport, DNA replication and apoptosis. GAPDH is a key enzyme in glycolysis that catalyzes the first step of the pathway by converting D-glyceraldehyde 3-phosphate (G3P) into 3-phospho-D-glyceroyl phosphate.

References

Note: This product is for in vitro research use only