UTF1 Antibody

Catalog No: #48109

SAB Signalway Antibody

Package Size: #48109-1 50ul #48109-2 100ul

Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


	മല	۱'n	\sim	\mathbf{a}	n
\boldsymbol{L}	esc	/I II	υu	v	
		١ ١			

se ocional 4	
4	
affinity purified	
ICC	
ls,Rt	
de	
1 antibody hUTF1 antibody Undifferentiated embryonic cell transcription factor 1 antibody UTF 1	
ody UTF1 antibody UTF1_HUMAN antibody	
s-Prot#:Q5T230	
230	
ç	
36-50 kDa	
1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.	
Store at -20°C	
1: c = :;	


Application Details

WB: 1:500-1:1000ICC: 1:500-1:1,000

Images



Western blot analysis on D3 cell lysates using anti- UTF1 mouse mAb.



ICC staining UTF1 in D3 cells (red). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Undifferentiated embryonic cell transcription factor 1 is a protein in humans that is encoded by the UTF1 gene. UTF1, first reported in 1998, is expressed in pluripotent cells including embryonic stem cells and embryonic carcinoma cells. Its expression is rapidly reduced upon differentiation. UTF1 protein is localized to the cell nucleus, where it functions to regulate the pluripotent chromatin state and buffer mRNA levels by promoting degradation of mRNA. Aberrant expression of UTF1 has also been reported in cervical cancer cells, where the UTF1 gene promoter loses methylation and becomes abnormally expressed compared to normal cervical cells.

References

Note: This product is for in vitro research use only