

CD137 Antibody

Catalog No: #48126

Package Size: #48126-1 50ul #48126-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	CD137 Antibody
Host Species	Mouse
Clonality	Monoclonal
Clone No.	2G1
Purification	ProA affinity purified
Applications	WB, IHC, ICC, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Recombinant protein
Other Names	4 1BB antibody 4 1BB ligand receptor antibody 4-1BB ligand receptor antibody 4-1BB Ligand Receptor T Cell antibody 4-1BB, mouse, homolog of antibody Antigen 4-1BB Homolog antibody CD 137 antibody CD137 antibody CD137 antigen antibody CDw137 antibody HLDA VI antibody Homolog of mouse 4 1BB antibody ILA antibody induced by lymphocyte activation (ILA) antibody Induced by lymphocyte activation antibody Interleukin activated receptor homolog of mouse Ly63 antibody Ly63, mouse, homolog of antibody MGC2172 antibody OTTHUMP00000044294 antibody Receptor protein 4 1BB antibody T cell antigen 4 1BB homolog antibody T cell antigen ILA antibody T-cell antigen 4-1BB homolog antibody T-cell antigen ILA antibody TNF receptor superfamily member 9 antibody TNFRSF9 antibody TNR9_HUMAN antibody Tumor necrosis factor receptor superfamily member 9 antibody
Accession No.	Swiss-Prot#:Q07011
Uniprot	Q07011
GeneID	3604;
Calculated MW	35 kDa
Formulation	1*TBS (pH7.4), 0.5%BSA, 50%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500 IHC: 1:50-1:200 ICC: 1:100-1:500FC: 1:50-1:200


Images




Western blot analysis of CD137 on HepG2 cell lysate using anti-CD137 antibody at 1/500 dilution.




Immunohistochemical analysis of paraffin-embedded rat brain tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded rat hippocampus tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue using anti-CD137 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-CD137 antibody. Counter stained with hematoxylin.




ICC staining CD137 (green) in A431 cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining CD137 (green) in A549 cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining CD137 (green) in LOVO cells. The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Jurkat cells with CD137 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black).

Background

CD137, also designated ILA and 4-1BB in mouse, belongs to the tumor necrosis factor receptor family and delivers a costimulatory signal to T lymphocytes. CD137 is expressed on activated T cells and binds an inducible ligand that is found on B cells, macrophages, and dendritic cells. Interactions between CD137 and its ligand are involved in antigen presentation and the generation of cytotoxic T cells. Crosslinking of the CD137 ligand induces apoptosis in resting lymphocytes. In contrast, CD137 regulates peripheral monocyte survival by inducing a cytokine release profile, and is mediated by M-CSF and to a lesser extent by granulocyte-macrophage colony-stimulating factor and IL-3. Soluble forms of CD137 are found in sera from patients with rheumatoid arthritis and may provide a negative control mechanism for immune responses.

References

Note: This product is for in vitro research use only