

## FAK (Phospho-Ser 722) Antibody

Catalog No: #13327

Package Size: #13327-1 50ul #13327-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

## Description

Product Name	FAK (Phospho-Ser 722) Antibody
Host Species	Mouse
Clone No.	4G1
Purification	ProA affinity purified
Applications	WB, IP, IF, IHC(P)
Species Reactivity	Hu, Ms, Rt
Immunogen Description	peptide
Other Names	FADK 1 antibody FADK antibody FAK related non kinase polypeptide antibody FAK1 antibody FAK1_HUMAN antibody Focal adhesion kinase 1 antibody Focal adhesion Kinase antibody Focal adhesion kinase isoform FAK Del33 antibody Focal adhesion kinase related nonkinase antibody FRNK antibody p125FAK antibody pp125FAK antibody PPP1R71 antibody Protein phosphatase 1 regulatory subunit 71 antibody Protein tyrosine kinase 2 antibody Protein-tyrosine kinase 2 antibody Ptk2 antibody PTK2 protein tyrosine kinase 2 antibody
Accession No.	Swiss-Prot#:Q05397
Uniprot	Q05397
GeneID	5747;
Calculated MW	125 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at 4°C


## Application Details

WB: 1:100-1:1,000


IHC: 1:50-1:500

IP: 1-2 &amp;mu;g per 100-500 &amp;mu;g of total protein(1 ml of cell lysate)

## Images



A. Western blot analysis of FAK phosphorylation in non-transfected:(A,D), untreated human FAK transfected:(B,E) and lambda protein phosphatase treated human FAK transfected:(C,F) 293T whole cell lysates. Antibodies tested include p-FAK (A-12):(A,B,C) and FAK (C-903):(D,E,F).



B. Immunoperoxidase staining of formalin fixed, paraffin-embedded human adrenal gland tissue showing cytoplasmic staining of glandular cells.

## Background

Activation of integrins in the extracellular matrix (ECM) of eukaryotic cells promotes the formation of membrane adhesion complexes, known as focal adhesions, which can include cytoskeletal proteins and protein tyrosine kinases, such as focal adhesion kinase (FAK). Phosphorylation events occurring within focal adhesions influence numerous processes that include mitogenic signaling, cell survival and cell motility. FAK is a non-receptor tyrosine kinase that is ubiquitously expressed and highly conserved between species. FAK is recruited by integrin clusters and variably phosphorylated depending on the effector molecules present in the focal adhesion. Phosphorylation of FAK Tyr 397 decreases during serum starvation, contact inhibition and cell cycle arrest, all conditions under which activating FAK Tyr 407 phosphorylation increases.

Note: This product is for in vitro research use only