xanthine dehydrogenase Antibody

Catalog No: #48373

Package Size: #48373-1 50ul #48373-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_				
	മഠ	cri	ntı	on
$\boldsymbol{ u}$	CO	OH	ν u	UH

Product Name	xanthine dehydrogenase Antibody		
Host Species	Mouse		
Clonality	Monoclonal		
Clone No.	1C8		
Purification	ProG affinity purified		
Applications	IHC,FC		
Species Reactivity	Hu		
Immunogen Description	Peptide		
Other Names	Xanthine dehydrogenase antibody Xanthine dehydrogenase/oxidase antibody Xanthine oxidase antibody		
	Xanthine oxidoreductase antibody XD antibody XDH antibody XDH_HUMAN antibody xdha antibody		
	XO antibody xor antibody		
Accession No.	Swiss-Prot#:P47989		
Uniprot	P47989		
GeneID	7498;		
Calculated MW	146 kDa		
Formulation	1*TBS (pH7.4), 0.5%BSA, 50%Glycerol. Preservative: 0.05% Sodium Azide.		
Storage	Store at -20°C		


Application Details

IHC: 1:50-1:200 FC: 1:50-1:100


Images



Immunohistochemical analysis of paraffin-embedded human liver tissue using anti-xanthine dehydrogenase antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-xanthine dehydrogenase antibody. Counter stained with hematoxylin.



Flow cytometric analysis of LOVO cells with xanthine dehydrogenase antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated Goat anti mouse IgG was used as the secondary antibody.

Background

The process of metabolizing purines to a common molecule known as xanthine is an essential process for the proper shuttling of uric acid. Xanthine oxidase is a flavoprotein enzyme that coordinates molybdenum and utilizes NAD+ as an electron acceptor to catalyze the oxidation of hypoxanthine to xanthine and then to uric acid. The predominant form of this enzyme is xanthine dehydrogenase, which is a homodimer that can be converted to xanthine oxidase by sulfhydryl oxidation or proteolytic modification. Xanthine oxidase is present in species ranging from bacteria to human and is ubiquitously expressed in mammalian tissues. In the oxidase form, this enzyme is coupled to the generation of free radicals. Individuals showing marked elevation of serum xanthine oxidase is suggestive of chronic liver disease and cholestasis, which is a condition defined by hepatic obstruction. Hepatic obstruction causes bile salts, the bile pigment bilirubin, and fats to accumulate in the blood stream instead of being eliminated normally. The clinical consequences of defects in xanthine oxidase range from mild to severe and even contribute to fatal disorders.

References

Note: This product is for in vitro research use only