DIS3L2 Antibody

Catalog No: #48435

Package Size: #48435-1 50ul #48435-2 100ul Orders:



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


$\overline{}$			
	escri	ntı	Λn
$\boldsymbol{ u}$	COUL	рu	OH

Product Name	DIS3L2 Antibody	
Host Species	Mouse	
Clonality	Monoclonal	
Clone No.	B2-C7	
Purification	ProA affinity purified	
Applications	WB, IHC, FC	
Species Reactivity	Hu	
Immunogen Description	Recombinant protein	
Other Names	DI3L2_HUMAN antibody DIS3 L2 antibody DIS3 mitotic control homolog (S. cerevisiae) like 2 antibody DIS3	
	mitotic control homolog like 2 antibody DIS3-like exonuclease 2 antibody Dis3l2 antibody FLJ36974 antibody	
	MGC42174 antibody	
Accession No.	Swiss-Prot#:Q8IYB7	
Uniprot	Q8IYB7	
GeneID	129563;	
Calculated MW	100 kDa	
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.	
Storage	Store at -20°C	


Application Details

WB: 1:500-1:1,000 IHC: 1:50-1:200FC: 1:100-1:200


Images




Western blot analysis of DIS3L2 on human DIS3L2 recombinant protein using anti-DIS3L2 antibody at 1/1,000 dilution.




Western blot analysis of DIS3L2 on HEK293 (1) and DIS3L2-hlgGFc transfected HEK293 (2) cell lysate using anti-DIS3L2 antibody at 1/1,000 dilution.




Western blot analysis of DIS3L2 on Hela (1) and HepG2 (2) cell lysate using anti-DIS3L2 antibody at 1/1,000 dilution.



Immunohistochemical analysis of paraffin-embedded human endometrial cancer tissue using anti-DIS3L2 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human bladder cancer tissue using anti-DIS3L2 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of Jurkat cells with DIS3L2 antibody at 1/100 dilution (green) compared with an unlabelled control (cells without incubation with primary antibody; red).

Background

The exosome is a multisubunit complex composed of several highly conserved subunits, some of which are 3' to 5' exoribonucleases. The complex is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich (ARE) elements in their untranslated 3' region. DIS3L2 (DIS3-like exonuclease 2) is an 885 amino acid protein that is thought to function as an exonuclease and may be required for the 3' processing of pre-mRNA into mature mRNA. Defects or chromosomal translocations involving the gene encoding DIS3L2 may be associated with Marfanoid habitus, a genetic disorder characterized by abnormalities in the skeleton, eyes and cardiovascular system. DIS3L2 is expressed as five isoforms due to alternative splicing events.

References

Note: This product is for in vitro research use only