

Phospho-NF-kB p65 (S529) Rabbit mAb

Catalog No: #13347



Package Size: #13347-1 50ul #13347-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	Phospho-NF-kB p65 (S529) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	SP07-00
Purification	ProA affinity purified
Applications	WB, IP, FC
Species Reactivity	Hu
Immunogen Description	Synthetic peptide conjugated to Keyhole Limpet Haemocyanin (KLH)
Other Names	Avian reticuloendotheliosis viral (v rel) oncogene homolog A antibody MGC131774 antibody NF kappa B p65delta3 antibody NFKB3 antibody Nuclear Factor NF Kappa B p65 Subunit antibody Nuclear factor NF-kappa-B p65 subunit antibody Nuclear factor of kappa light polypeptide gene enhancer in B cells 3 antibody Nuclear factor of kappa light polypeptide gene enhancer in B-cells 3 antibody OTTHUMP00000233473 antibody OTTHUMP00000233474 antibody OTTHUMP00000233475 antibody OTTHUMP00000233476 antibody OTTHUMP00000233900 antibody p65 antibody p65 NF kappaB antibody p65 NFkB antibody relA antibody TF65_HUMAN antibody Transcription factor p65 antibody v rel avian reticuloendotheliosis viral oncogene homolog A (nuclear factor of kappa light polypeptide gene enhancer in B cells 3 (p65)) antibody V rel avian reticuloendotheliosis viral oncogene homolog A antibody v rel reticuloendotheliosis viral oncogene homolog A (avian) antibody V rel reticuloendotheliosis viral oncogene homolog A, nuclear factor of kappa light polypeptide gene enhancer in B cells 3, p65 antibody
Accession No.	Swiss-Prot#:Q04206
Uniprot	Q04206
GeneID	5970;
Calculated MW	60 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:1,000 FC: 1:50-1:100

Images



Flow cytometric analysis of Daudi cells with Phospho-NF-kB p65(S529) antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Proteins encoded by the *v-Rel* viral oncogene and its cellular homolog, *c-Rel*, are members of a family of transcription factors that include the two subunits of the transcription factor $\text{NF-}\kappa\text{B}$ (p50 and p65) and the *Drosophila* maternal morphogen, dorsal. Both proteins specifically bind to DNA sequences that are the same or slight variations of the 10 bp κB sequence in the immunoglobulin κ light chain enhancer. This same sequence is also present in a number of other cellular and viral enhancers. The DNA binding activity of $\text{NF-}\kappa\text{B}$ is activated and $\text{NF-}\kappa\text{B}$ is subsequently transported from the cytoplasm to the nucleus in cells exposed to mitogens or growth factors. cDNAs encoding precursors for two distinct proteins of the same size have been described, designated p105 and p100. The p105 precursor contains p50 at its N-terminus and a C-terminal region that when expressed as a separate molecule, designated p Δ 1, binds to p50 and regulates its activity.

References

Note: This product is for in vitro research use only