

CD86 Rabbit mAb

Catalog No: #48763

Package Size: #48763-1 50ul #48763-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

| | |
|-----------------------|--|
| Product Name | CD86 Rabbit mAb |
| Clone No. | SJ20-00 |
| Purification | ProA affinity purified |
| Applications | WB, ICC/IF, IHC, IP, FC |
| Species Reactivity | Hu, Rt |
| Immunogen Description | Synthetic peptide within Human CD86 aa 1-50 / 329. |
| Other Names | Activation B7-2 antigen 3 antibody Activation B7-2 antigen antibody B-lymphocyte activation antigen B7-2 2 antibody B-lymphocyte activation antigen B7-2 antibody B7 2 antibody B7 antibody B7-2 antibody B7.2 antibody B70 antibody B72 antigen antibody BU63 antibody CD28 antigen ligand 2 2 antibody CD28 antigen ligand 2 antibody Cd28l2 antibody CD28LG2 antibody CD86 antibody CD86 antigen (CD28 antigen ligand 2, B7-2 antigen) 1, 2 antibody CD86 antigen (CD28 antigen ligand 2, B7-2 antigen) antibody CD86 antigen antibody CD86 molecule antibody CD86_HUMAN antibody CLS1 antibody CTLA-4 counter-receptor B7.2 2, 3 antibody CTLA-4 counter-receptor B7.2 antibody Early T-cell costimulatory molecule 1 antibody ETC-1 antibody FUN 1 antibody FUN-1 antibody LAB72 antibody Ly-58 antibody Ly58 antibody MB7 antibody MB7-2 antibody Membrane glycoprotein antibody MGC34413 antibody T-lymphocyte activation antigen CD86 antibody TS/A-2 antibody |
| Accession No. | Swiss-Prot#:P42081 |
| Uniprot | P42081 |
| GeneID | 942; |
| Calculated MW | 38 kDa |
| Concentration | 1 mg/ml |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |

Application Details

WB: 1:500-1:2,000


IHC: 1:50-1:200

ICC: 1:50-1:200


FC: 1:50-1:100

IP: Use at an assay dependent concentration.


Images




Western blot analysis of CD86 on different lysates using anti-CD86 antibody at 1/1,000 dilution. Positive control:
Lane 1: Raji
Lane 2: Jurkat




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-CD86 antibody. Counter stained with hematoxylin.




ICC staining CD86 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining CD86 in HUVEC cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining CD86 in JAR cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of K562 cells with CD86 antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

T cell proliferation and lymphokine production are triggered by occupation of the TCR by antigen, followed by a costimulatory signal that is delivered by a ligand expressed on antigen presenting cells. The B7-related cell surface proteins B7-1 (CD80) and B7-2 (CD86) expressed on antigen presenting cells bind the homologous T cell receptors CD28 and CTLA-4 (cytotoxic T lymphocyte-associated protein-4) and trigger costimulatory signals for optimal T cell activation. CTLA-4 shares 31% overall amino acid identity with CD28, and it has been proposed that CD28 and CTLA-4 are functionally redundant. SLAM is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. B7, also designated BB1, is another ligand or counterreceptor for CD28 and CTLA-4 that is expressed on the antigen-presenting cell.

Note: This product is for in vitro research use only