Phospho-JNK1/2/3(T183+T183+T221) Rabbit mAb

Catalog No: #13371

SAB Signalway Antibody

Package Size: #13371-1 50ul #13371-2 100ul


Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Product Name	Phospho-JNK1/2/3(T183+T183+T221) Rabbit mAb
Host Species	Rabbit
Clonality	Monoclonal
Clone No.	ST500
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu, Ms, Rt
mmunogen Description	Synthetic phospho-peptide corresponding to residues surrounding Thr183 + Thr183 + Thr221 of human JNK1/2/3
Other Names	C Jun kinase 2 antibody c Jun N terminal kinase 1 antibody c Jun N terminal kinase 2 antibody c Jun N terminal kinase 3 antibody c Jun N terminal kinase 3 antibody c Jun N terminal kinase 1 antibody JNK 46 antibody JNK 55 antibody JNK antibody JNK-46 antibody JNK-48 antibody JNK-48 antibody JNK-48 antibody JNK-48 antibody JNK-48 antibody JNK-48 antibody MAP kinase 9 antibody MAP kinase 9 antibody MAP kinase 9 antibody MAP kinase 9 antibody MAP kinase 8 isoform JNK-1 alpha-1 antibody Mitogen activated protein kinase 8 isoform JNK-1 beta-2 antibody Mitogen activated protein kinase 9 antibody Mitogen activated protein kinase 8 antibody MK-48 antibody PRK-48 antibod
Accession No.	Swiss-Prot#:P45983
Jniprot	P45983
GeneID	5599;
Calculated MW	54/46 kDa
Saloulatoa IIIII	

Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100

Images




Western blot analysis of Phospho-JNK1/2/3(T183+T183+T221) on different lysates using anti-Phospho-JNK1/2/3(T183+T183+T221) antibody at


Lane 1: NIH/3T3 cell lysate, treated with Anisomycin

Lane 2: NIH/3T3 cell lysate, untreated


1/1,000 dilution. Positive control:




Immunohistochemical analysis of paraffin-embedded human liver tissue using anti-Phospho-JNK1/2/3(T183+T183+T221) antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human uterus tissue using anti-Phospho-JNK1/2/3(T183+T183+T221) antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue using anti-Phospho-JNK1/2/3(T183+T183+T221) antibody. Counter stained with hematoxylin.




ICC staining Phospho-JNK1/2/3(T183+T183+T221) in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Phospho-JNK1/2/3(T183+T183+T221) in NIH/3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Phospho-JNK1/2/3(T183+T183+T221) in HUVEC cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Hela cells with Phospho-JNK1/2/3(T183+T183+T221) antibody at 1/50 dilution (blue) compared with an unlabelled control (cells without incubation with primary antibody; red). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

JNKs (c-Jun N-terminal kinases) belong to a family of MAP kinases that are involved in a variety of cellular processes, including transcriptional regulation and cellular proliferation, differentiation and development. JNK2 (c-Jun N-terminal kinase 2) and JNK3 (c-Jun N-terminal kinase 3) are 424 and 464 amino acid proteins, respectively, that each contain one protein kinase domain and use magnesium as a cofactor to catalyze the phosphorylation of target proteins, thereby playing a role in a variety of events throughout the cell. Both JNK2 and JNK3 exist as multiple alternatively spliced isoforms and are subject to post-translational phosphorylation on Thr 183 and Thr 221, respectively, an event which activates JNK2/JNK3 enzymatic activity. Defects in the gene encoding JNK3 are a cause of epileptic encephalopathy of the Lennox-Gastaut type, a group of epileptic disorders characterized by severe psychomotor delay and seizures.

References

Note: This product is for in vitro research use only