

Moesin Rabbit mAb

Catalog No: #48981

Package Size: #48981-1 50ul #48981-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

| | |
|-----------------------|--|
| Product Name | Moesin Rabbit mAb |
| Host Species | Recombinant Rabbit |
| Clonality | Monoclonal antibody |
| Clone No. | SC69-01 |
| Purification | ProA affinity purified |
| Applications | WB, ICC/IF, IHC, IP, FC |
| Species Reactivity | Hu, Ms, Rt |
| Immunogen Description | recombinant protein |
| Other Names | Epididymis luminal protein 70 antibody HEL70 antibody Membrane organizing extension spike protein antibody Membrane-organizing extension spike protein antibody MOES_HUMAN antibody Moesin antibody Moesin/anaplastic lymphoma kinase fusion protein antibody Msn antibody MSN/ALK fusion antibody |
| Accession No. | Swiss-Prot#:P26038 |
| Uniprot | P26038 |
| GeneID | 4478; |
| Calculated MW | 68 kDa |
| Formulation | 1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide. |
| Storage | Store at -20°C |


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


Images




Western blot analysis of Moesin on different lysates using anti-Moesin antibody at 1/1,000 dilution. Positive control:
Lane 1: Raji
Lane 2: SH-SY-5Y




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Moesin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-Moesin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-Moesin antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded mouse lung tissue using anti-Moesin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse placenta tissue using anti-Moesin antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse stomach tissue using anti-Moesin antibody. Counter stained with hematoxylin.



ICC staining Moesin in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Moesin in NCCIT cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of HeLa cells with Moesin antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Ezrin, Moesin and Radixin belong to a family of highly homologous Actin-associated proteins that are localized just beneath the plasma membrane. These proteins are believed to be involved in the mediation of interactions between cytoskeletal and membrane proteins. Ezrin serves as a major cytoplasmic substrate of various protein-tyrosine kinases, including the epidermal growth factor receptor. Ezrin has also been identified as a cAMP-dependent protein kinase (A-kinase) anchoring protein and designated AKAP78. Moesin and Radixin share more than 70% homology with Ezrin and are co-expressed within various cell types. Despite the high degree of homology, the three proteins exhibit a distinct receptor-specific pattern of phosphorylation.

References

Note: This product is for in vitro research use only