

PYK2 Rabbit mAb

Catalog No: #49000

Package Size: #49000-1 50ul #49000-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	PYK2 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SC06-15
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Other Names	CADTK antibody CAK-beta antibody CAKB antibody CAKbeta antibody Calcium regulated non receptor proline rich tyrosine kinase antibody Calcium-dependent tyrosine kinase antibody Cell adhesion kinase beta antibody E430023O05Rik antibody EC 2.7.10.2 antibody FADK 2 antibody FADK2 antibody FAK2 antibody FAK2_HUMAN antibody Focal adhesion kinase 2 antibody MGC124628 antibody PKB antibody Proline-rich tyrosine kinase 2 antibody Protein kinase B antibody Protein Tyrosine Kinase 2 Beta antibody Protein-tyrosine kinase 2-beta antibody PTK antibody PTK2B antibody PTK2B protein tyrosine kinase 2 beta antibody PYK2 antibody RAFTK antibody RAFTK2 antibody Related adhesion focal tyrosine kinase antibody
Accession No.	Swiss-Prot#:Q14289
Uniprot	Q14289
GeneID	2185;
Calculated MW	116 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200


Images




Western blot analysis of PYK2 on mouse brain lysates using anti-PYK2 antibody at 1/1,000 dilution.




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-PYK2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-PYK2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-PYK2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse brain tissue using anti-PYK2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse kidney tissue using anti-PYK2 antibody. Counter stained with hematoxylin.



ICC staining PYK2 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining PYK2 in SHG-44 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining PYK2 in SH-SY-5Y cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Focal adhesion kinase (FAK) was initially identified as a substrate for the intrinsic protein tyrosine kinase activity of Src-encoded pp60. The deduced amino acid sequence of FAK p125 has shown it to be a cytoplasmic protein tyrosine kinase whose sequence and structural organization are unique compared to other protein families described. A putative new member of the FAK family, designated PYK2 (proline-rich tyrosine kinase 2), exhibits 61% sequence identity with FAK over its kinase domain. PYK2 (also designated CAKb or RAFTK) is highly expressed in the central nervous system. Activation of the kinase leads to modulation of ion channel function and the activation of the MAPK signaling pathway. PYK2 is rapidly phosphorylated on tyrosine residues in response to stimuli that increase intracellular calcium levels and compounds that activate members of the PKC family of kinases, such as phorbol esters.

References

Note: This product is for in vitro research use only