

IRF7 Rabbit mAb

Catalog No: #49007

Package Size: #49007-1 50ul #49007-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	IRF7 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SC0617
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu, Ms, Rt, zebrafish
Immunogen Description	recombinant protein
Other Names	IMD39 antibody Interferon regulatory factor 7 antibody Interferon regulatory factor 7H antibody IRF 7 antibody IRF 7A antibody IRF 7H antibody IRF-7 antibody IRF7 antibody IRF7_HUMAN antibody IRF7A antibody IRF7B antibody IRF7C antibody IRF7H antibody
Accession No.	Swiss-Prot#:Q92985
Uniprot	Q92985
GeneID	3665;
Calculated MW	54 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:100-1:500FC: 1:50-1:100


Images




Western blot analysis of IRF7 on different lysates using anti-IRF7 antibody at 1/1,000 dilution. Positive control:
Lane 1: Jurkat
Lane 2: Raji




Western blot analysis of IRF7 on hybrid fish (crucian-carp) brain tissue lysate using anti-IRF7 antibody at 1/500 dilution.




Immunohistochemical analysis of paraffin-embedded mouse heart tissue using anti-IRF7 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-IRF7 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human spleen tissue using anti-IRF7 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-IRF7 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue using anti-IRF7 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded mouse kidney tissue using anti-IRF7 antibody. Counter stained with hematoxylin.



ICC staining IRF7 in PC-12 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining IRF7 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding factors that function as regulators of both type I interferon (interferon- α and β) and interferon-inducible genes. The two factors are structurally related, particularly in their N-terminal regions, which confer DNA binding specificity. In addition, both bind to the same sequence within the promoters of interferon- α and interferon- β genes. IRF-1 functions as an activator of interferon transcription, while IRF-2 binds to the same cis elements and represses IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in regulating cell growth; overexpression of the repressor IRF-2 leads to cell transformation while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7 and IFN consensus sequence-binding protein (ICSBP).

References

Note: This product is for in vitro research use only