

Cyclin E1 Rabbit mAb

Catalog No: #49124

Package Size: #49124-1 50ul #49124-2 100ul

Orders: order@signalwayantibody.comSupport: tech@signalwayantibody.com


Description

Product Name	Cyclin E1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SD20-24
Purification	ProA affinity purified
Applications	WB, ICC/IF
Species Reactivity	Hu
Immunogen Description	recombinant protein
Other Names	CCNE antibody Ccne1 antibody CCNE1_HUMAN antibody cyclin E variant ex5del antibody cyclin E variant ex7del antibody Cyclin E1 antibody Cyclin Es antibody Cyclin Et antibody CyclinE antibody G1/S specific cyclin E antibody G1/S-specific cyclin-E1 antibody
Accession No.	Swiss-Prot#:P24864
Uniprot	P24864
GeneID	898;
Calculated MW	48 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000 ICC: 1:50-1:200


Images



ICC staining Cyclin E1 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Cyclin E1 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Cyclin E1 in MCF-7 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

Background

Cyclins were first identified in invertebrates as proteins that oscillate dramatically through the cell cycle. These proteins have been well conserved through evolution and play a critical role in regulation of cell division. cyclin E, along with the three cyclin D proteins and cyclin C, has been shown to represent a putative G1 cyclin on the basis of its cyclic pattern of mRNA expression, with maximal levels being detected near the G1/S boundary. cyclin E has been found to be associated with the transcription factor E2F in a temporally regulated manner. The cyclin E/E2F complex is detected primarily during the G1 phase of the cell cycle and decreases as cells enter S phase. E2F is known to be a critical transcription factor for expression of several S phase specific proteins.

References

Note: This product is for in vitro research use only