

Cdk4 Rabbit mAb

Catalog No: #49132

Package Size: #49132-1 50ul #49132-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description


Product Name	Cdk4 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SD205-1
Purification	ProA affinity purified
Applications	WB, ICC/IF, IP
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Other Names	Cdk 4 antibody cdk4 antibody CDK4 protein antibody CDK4_HUMAN antibody Cell division kinase 4 antibody Cell division protein kinase 4 antibody CMM 3 antibody CMM3 antibody Crk3 antibody Cyclin dependent kinase 4 antibody Cyclin-dependent kinase 4 antibody Melanoma cutaneous malignant 3 antibody MGC14458 antibody p34 cdk4 antibody PSK J3 antibody PSK-J3 antibody
Accession No.	Swiss-Prot#:P11802
Uniprot	P11802
GeneID	1019;
Calculated MW	34 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details


WB: 1:1,000-1:2,000

ICC: 1:50-1:200


Images




Western blot analysis of Cdk4 on different lysates using anti-Cdk4 antibody at 1/1,000 dilution. Positive control: Lane 1: Hela Lane 2: MCF-7 Lane 3: K562




ICC staining Cdk4 in HeLa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Cdk4 in AGS cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining Cdk4 in NIH/3T3 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Western blotting analysis using Cdk4 Antibody #49132.

Background

Cell cycle progression is controlled in part by a family of cyclin proteins and cyclin dependent kinases (Cdks). Cdk proteins work in concert with the cyclins to phosphorylate key substrates involved in each phase of cell cycle progression. Another family of proteins, Cdk inhibitors, also plays a role in regulating the cell cycle by binding to cyclin-Cdk complexes and modulating their activity. Several Cdk proteins have been identified, including Cdk2-Cdk8, PCTAIRE-1-PCTAIRE-3, PITALRE and PITSLRE. Cdk4, in complex with D-type cyclins, is thought to regulate cell growth during the G1 phase of the cell cycle. This association with a D-type cyclin upregulates Cdk4 activity, whereas binding to the Cdk inhibitor p16 downregulates Cdk4 activity. Activation of the Cdk4-cyclin complexes requires phosphorylation on a single threonyl residue of Cdk4, catalyzed by a Cdk-activating protein (CAK).

Note: This product is for in vitro research use only