

Histone H2B Rabbit mAb

Catalog No: #49134



Package Size: #49134-1 50ul #49134-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


Description

Product Name	Histone H2B Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	SD20-63
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC, IP, FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	recombinant protein
Other Names	H2B GL105 antibody H2B histone family member O antibody H2B histone family member S antibody H2B.1 antibody H2B.1 B antibody H2B.b antibody H2B.c antibody H2B.d antibody H2B.e antibody H2B.f antibody H2B.j antibody H2B.q antibody H2B/b antibody H2B/c antibody H2B/d antibody H2B/e antibody H2B/f antibody H2B/j antibody H2B/o antibody H2B/q antibody H2BFB antibody H2BFC antibody H2BFD antibody H2BFE antibody H2BFF antibody H2BFJ antibody H2BFO antibody H2BFQ antibody H2BFS antibody HIRIP2 antibody HIST1H2BB antibody HIST1H2BD antibody HIST1H2BH antibody HIST1H2BL antibody HIST1H2BM antibody HIST1H2BN antibody HIST2H2BE antibody Histone H2B antibody Histone H2B type 1 B antibody Histone H2B type 1 D antibody Histone H2B type 1 H antibody Histone H2B type 1 L antibody Histone H2B type 1 M antibody Histone H2B type 1 N antibody Histone H2B type 2 E antibody histone protein antibody
Accession No.	Swiss-Prot#:O60814
Uniprot	O60814
GeneID	85236;
Calculated MW	14 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:1,000-5,000 IHC: 1:50-1:200 ICC: 1:100-1:500 FC: 1:50-1:100


Images




Western blot analysis of Histone H2B on different lysates using anti-Histone H2B antibody at 1/1,000 dilution. Positive control: Lane 1: Hela Lane 2: NIH/3T3 Lane 3: PC12




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Histone H2B antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human liver tissue using anti-Histone H2B antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse liver tissue using anti-Histone H2B antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse testis tissue using anti-Histone H2B antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse colon tissue using anti-Histone H2B antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-Histone H2B antibody. Counter stained with hematoxylin.



ICC staining Histone H2B in A431 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of HeLa cells with Histone H2B antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

Eukaryotic histones are basic and water soluble nuclear proteins that form hetero-octameric nucleosome particles by wrapping 146 base pairs of DNA in a left-handed super-helical turn sequentially to form chromosomal fiber. Two molecules of each of the four core histones (H2A, H2B, H3, and H4) form the octamer; formed of two H2A-H2B dimers and two H3-H4 dimers, forming two nearly symmetrical halves by tertiary structure. Over 80% of nucleosomes contain the linker Histone H1, derived from an intronless gene, that interacts with linker DNA between nucleosomes and mediates compaction into higher order chromatin. Histones are subject to posttranslational modification by enzymes primarily on their N-terminal tails, but also in their globular domains. Such modifications include methylation, citrullination, acetylation, phosphorylation, sumoylation, ubiquitination and ADP-ribosylation.

References

Note: This product is for in vitro research use only