CD3D Rabbit mAb

Catalog No: #49268

Package Size: #49268-1 50ul #49268-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

CD3 antigen delta subunit antibody CD3 delta antibody CD3d antibody CD3d antigen delta polypeptide (TiT3			
complex) antibody CD3d molecule delta (CD3-TCR complex) antibody CD3D_HUMAN antibody IMD19			
antibody OKT3 delta chain antibody T cell receptor T3 delta chain antibody T-cell receptor T3 delta chain			
ly			
hloride, 0.05% BSA, 0.02% sodium			
w cycle.			


Application Details

IHC 1:100-1:200 ICC/IF 1:50-1:200 IP 1:20-1:50	WB 1:1000-1:2000			
	IHC 1:100-1:200			
IP 1:20-1:50	ICC/IF 1:50-1:200			
	IP 1:20-1:50			
FC 1:20-1:100	FC 1:20-1:100			


Images




Flow cytometric analysis of Raji cells with CD3D antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.




All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



All lanes use the Antibody at 1:2K dilution for 1 hour at room temperature.



Immunohistochemical analysis of paraffin-embedded human spleen, using CD3D Antibody.



Immunofluorescent analysis using the Antibody at 1:50 dilution.

8

Immunofluorescent analysis of Jurkat cells, using CD3D Antibody.

Part of the TCR-CD3 complex present on T-lymphocyte cell surface that plays an essential role in adaptive immune response.

Note: This product is for in vitro research use only