

## IGF2 Rabbit mAb

Catalog No: #49302

Package Size: #49302-1 50ul #49302-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	IGF2 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JJ092-3
Purification	ProA affinity purified
Applications	WB, ICC/IF, IHC
Species Reactivity	Hu
Immunogen Description	recombinant protein
Other Names	C11orf43 antibody IGF 2 antibody IGF II antibody IGF-II antibody IGF2 antibody IGF2_HUMAN antibody IGFII antibody INSIGF antibody insulin like growth factor 2 (somatomedin A) antibody Insulin like Growth Factor 2 antibody Insulin like growth factor II antibody Insulin like growth factor II precursor antibody Insulin like growth factor type 2 antibody pp9974 antibody Preptin antibody putative insulin like growth factor II associated protein antibody Somatomedin A antibody Somatomedin-A antibody
Accession No.	Swiss-Prot#:P01344
Uniprot	P01344
GeneID	3481;
Calculated MW	11 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:1,000 IHC: 1:50-1:200 ICC: 1:100-1:500


## Images




Western blot analysis of IGF2 on human placenta lysates using anti-IGF2 antibody at 1/1,000 dilution.




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue using anti-IGF2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue using anti-IGF2 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human breast carcinoma tissue using anti-IGF2 antibody. Counter stained with hematoxylin.




ICC staining IGF2 in Hela cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining IGF2 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining IGF2 in 293T cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining IGF2 in 293 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.

## Background

IGF-II (Insulin-like growth factor II; also multiplication-stimulating polypeptide/MSP and somatomedin-A) is a secreted 8 kDa polypeptide that belongs to the insulin family of peptide growth factors. It is part of a complex system of growth and metabolic-regulating proteins that is particularly important during development. It has been associated with nervous system proliferation and differentiation, myelination, adrenal cortical proliferation, and skeletal growth and differentiation. In human, IGF-II is primarily synthesized by the liver, and circulates at high levels in both fetus and adult. In rodent, however, IGF-II levels drop after the perinatal period, an effect attributed to the lack of a key gene promoter. This may indicate that postnatally, IGF-II has either a limited, or local effect only in rodent.

## References

Note: This product is for in vitro research use only