SMC3 Rabbit mAb

Catalog No: #49448

Package Size: #49448-1 50ul #49448-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description					
Product Name	SMC3 Rabbit mAb				
Host Species	Recombinant Rabbit				
Clonality	Monoclonal antibody				
Clone No.	JM10-75				
Purification	ProA affinity purified				
Applications	WB, ICC/IF, IHC, FC				
Species Reactivity	Hu, Ms, Rt				
mmunogen Description	recombinant protein				
Other Names	BAM antibody Bamacan antibody Basement membrane associated chondroitin proteoglycan antibody				
	Basement membrane-associated chondroitin proteoglycan antibody BMH antibody CDLS3 antibody				
	chondroitin sulfate proteoglycan 6 (bamacan) antibody Chondroitin sulfate proteoglycan 6 antibody				
	Chromosome associated polypeptide antibody Chromosome-associated polypeptide antibody CSPG 6				
	antibody CSPG6 antibody hCAP antibody Human chromosome associated polypeptide antibody				
	im:7142991 antibody SMC 3 antibody SMC protein 3 antibody SMC-3 antibody smc3 antibody				
	SMC3_HUMAN antibody SMC3L1 antibody Structural maintenance of chromosome 3 antibody Structura				
	maintenance of chromosomes 3 antibody Structural maintenance of chromosomes protein 3 antibody				
	u:fb22e01 antibody wu:fc30d07 antibody				
Accession No.	Swiss-Prot#:Q9UQE7				
Jniprot	Q9UQE7				
GeneID	9126;				
Calculated MW	141 kDa				
ormulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.				
Storage	Store at -20°C				

Application Details


WB: 1:1,000-5,000IHC: 1:50-1:200 ICC: 1:100-1:500FC: 1:50-1:100

Images




Western blot analysis of SMC3 on different cells lysates using anti-SMC3 antibody at 1/1,000 dilution. Positive control:


Line1: HepG2 Line2: NIH-3T3 Line3:PC12




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-SMC3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human colon cancer tissue using anti-SMC3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded mouse colon tissue using anti-SMC3 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded zebrafish tissue using anti-SMC3 antibody. Counter stained with hematoxylin.




ICC staining SMC3 in Hela cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining SMC3 in HepG2 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining SMC3 in NIH-3T3 cells (red). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of Hela cells with SMC3 antibody at 1/50 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti rabbit IgG was used as the secondary antibody.

Background

The SMC (structural maintenance of chromosomes) family of proteins form heterodimeric complexes that modulate sister chromatid cohesion and chromosome condensation for mitosis. The two distinct classes of SMC protein complexes are comprised of SMC1 (also designated SB1.8) with SMC3 (also designated HCAP for human chromosome-associated protein and Bamacan for the secreted proteoglycan), and SMC2 (also designated hCAP-E) with SMC4 (also designated hCAP-C). The SMC1/SMC3 complex is required for metaphase progression in mitotic cells and functions independently of the SMC2/SMC4 complex during the cell cycle. SMC1 is ubiqitiously expressed in various human tissues, including thymus, testis, and colon. SMC3 is expressed as a nuclear protein in the colon, but can also occur as a secreted proteoglycan expressed in testis and brain. The secreted proteoglycan contains several glycosylation sites and is thought to play a role in basement membrane physiology.

_				
\mathbf{I}	\sim 4	-	•	ces
ĸ			n	\sim

Note: This product is for in vitro research use only