Hepcidin Rabbit mAb

Catalog No: #49534

Package Size: #49534-1 50ul #49534-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

n	000	rin	tio	n
	ヒゝし	IU	лю	

Hepcidin Rabbit mAb		
Recombinant Rabbit		
Monoclonal antibody		
JA11-27		
ProA affinity purified		
WB, IHC		
eactivity Hu, Rt		
recombinant protein		
Hamp antibody HEPC antibody HEPC_HUMAN antibody Hepc20 antibody Hepc25 antibody Hepcidin-20		
antibody HFE2B antibody LEAP-1 antibody LEAP1 antibody Liver-expressed antimicrobial peptide 1		
antibody PLTR antibody Putative liver tumor regressor antibody		
Swiss-Prot#:P81172		
P81172		
57817;		
3 kDa		
1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.		
Store at -20°C		

Application Details

WB: 1:500-1:1,000 IHC: 1:50-1:200

Images



Immunohistochemical analysis of paraffin-embedded human liver tissue using anti- Hepcidin antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded rat heart tissue using anti- Hepcidin antibody. Counter stained with hematoxylin.

Immunohistochemical analysis of paraffin-embedded rat brain tissue using anti- Hepcidin antibody. Counter stained with hematoxylin.

Background

Hepcidin, also known as HAMP, HEPC, LEAP1 or HFE2B, is an 84 amino acid secreted protein that regulates iron-related signaling events. Highly expressed in liver with lower expression in heart, brain, lung, prostate and thyroid, hepcidin is thought to maintain iron homeostasis and, in conjunction with the HFE protein (a protein that is defective in hereditary hemochromatosis), may mediate both iron storage in macrophages and intestinal iron absorption. Additionally, hepcidin has strong antimicrobial activity against gram-positive and gram-negative bacteria, as well as certain yeast strains, suggesting that hepcidin may play a crucial role in staving off bacterial infections. Defects in the gene encoding hepcidin are the cause of hemochromatosis type 2B (also known as juvenile hemochromatosis), an early-onset autosomal recessive disorder that results in severe iron overload and is characterized by hepatic fibrosis, hypogonadotrophic hypogonadism and cardiomyopathy.

References

Note: This product is for in vitro research use only