Eg5 Rabbit mAb

Catalog No: #49691

Package Size: #49691-1 50ul #49691-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Eg5 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JU83-16
Purification	ProA affinity purified
Applications	WB,IP
Species Reactivity	Hu
Immunogen Description	Recombinant protein
Other Names	EG5 antibody HKSP antibody KIF11 antibody KIF11_HUMAN antibody Kinesin family member 11 antibody
	Kinesin like protein 1 antibody Kinesin-like protein 1 antibody Kinesin-like protein KIF11 antibody Kinesin-like
	spindle protein HKSP antibody Kinesin-related motor protein Eg5 antibody KNSL1 antibody MCLMR
	antibody Thyroid receptor-interacting protein 5 antibody TR-interacting protein 5 antibody TRIP-5 antibody
	TRIP5 antibody
Accession No.	Swiss-Prot#:P52732
Uniprot	P52732
GenelD	3832;
Calculated MW	119 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C

Application Details

WB: 1:500-1:2,000IP: 1:10-1:50

Images



Background

Eukaryotes contain a superfamily of microtubule-based motor proteins comprising kinesin and a number of related proteins that are thought to participate in various forms of intracellular motility, including cell division and organelle transport. KIF11(also known as kinesin family member 11, Eg5 or TRIP5) is a slow, plus-end-directed microtubule-based motor of the BimC kinesin family that is essential for bipolar spindle formation during eukaryotic cell division. When the expression of KIF11 is blocked, centrosome migration halts and cells are arrested in mitosis with monoastral microtubule arrays. KIF11 is phosphorylated on serine during S phase and on both serine and Thr 927 during mitosis, which regulates the association of Eg5 with the spindle apparatus (probably during early prophase). KIF11 is also known to be a member of the thyroid receptor interacting protein (Trip) family, and interacts with the thyroid hormone receptor only in the presence of thyroid hormone.

References

Note: This product is for in vitro research use only