Ctip1 Rabbit mAb

Catalog No: #49820

Package Size: #49820-1 50ul #49820-2 100ul




Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Ctip1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JB35-68
Purification	ProA affinity purified
Applications	WB,IHC,FC
Species Reactivity	Hu
Immunogen Description	Recombinant protein
Other Names	2810047E18Rik antibody B cell CLL/lymphoma 11A (zinc finger protein) antibody B cell CLL/lymphoma
	11A (zinc finger protein) isoform 2 antibody B-cell CLL/lymphoma 11A antibody B-cell lymphoma/leukemia
	11A antibody BC11A_HUMAN antibody BCL-11A antibody BCL11A antibody BCL11A B cell
	CLL/lymphoma 11A (zinc finger protein) isoform 1 antibody BCL11A L antibody BCL11A S antibody
	BCL11A XL antibody BCL11a-M antibody BCL11AL antibody BCL11AS antibody BCL11AXL
	antibody C2H2 type zinc finger protein antibody COUP TF interacting protein 1 antibody
	COUP-TF-interacting protein 1 antibody CTIP1 antibody CTIP1, mouse, homolog of antibody
	D930021L15Rik antibody Ecotropic viral integration site 9 antibody Ecotropic viral integration site 9
	homolog antibody Ecotropic viral integration site 9 protein antibody Ecotropic viral integration site 9
	protein homolog antibody EVI-9 antibody Evi9 antibody Evi9, mouse, homolog of antibody FLJ10173
	antibody FLJ34997 antibody HBFQTL5 antibody KIAA1809 antibody mKIAA1809 antibody
	OTTHUMP00000159788 antibody OTTHUMP00000159789 antibody OTTHUMP00000201250 antibody
	OTTHUMP00000202084 antibody Zinc finger protein 856 antibody ZNF856 antibody
Accession No.	Swiss-Prot#:Q9H165
Uniprot	Q9H165
GenelD	53335;
Calculated MW	91 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:500-1:1,000 IHC: 1:50-1:200 FC: 1:50-1:100


## Images




Western blot analysis of Ctip1 on Daudi (1) and U937 (2) cell lysate using anti-Ctip1 antibody at 1/500 dilution.



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-Ctip1 antibody. Counter stained with hematoxylin.



Immunohistochemical analysis of paraffin-embedded mouse cerebellum tissue using anti-Ctip1 antibody. Counter stained with hematoxylin.



Flow cytometric analysis of Jurkat cells with Ctip1 antibody at 1/100 dilution (red) compared with an unlabelled control (cells without incubation with primary antibody; black). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

## Background

Transcription factor associated with the BAF SWI/SNF chromatin remodeling complex. Repressor of fetal hemoglobin (HbF) level. Involved in brain development. Functions as a myeloid and B-cell proto-oncogene. May play important roles in leukemogenesis and hematopoiesis. Essential factor in lymphopoiesis required for B-cell formation in fetal liver. May function as a modulator of the transcriptional repression activity of ARP1.

## References

Note: This product is for in vitro research use only