

ELMO1 Rabbit mAb

Catalog No: #49881



Package Size: #49881-1 50ul #49881-2 100ul

Orders: order@signalwayantibody.com
Support: tech@signalwayantibody.com


Description

Product Name	ELMO1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JG34-59
Purification	ProA affinity purified
Applications	WB,IHC,FC,IP
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Recombinant protein within human ELMO1 aa 500 to 700.
Other Names	CED 12 antibody Ced 12 homolog 1 antibody Ced 12 homolog antibody CED-12 antibody CED12 antibody Ced12 homolog 1 antibody Ced12 homolog antibody ELMO 1 antibody ELMO-1 antibody Elmo1 antibody ELMO1_HUMAN antibody Engulfment and cell motility 1 antibody Engulfment and cell motility protein 1 antibody KIAA0281 antibody MGC126406 antibody Protein ced-12 homolog antibody
Accession No.	Swiss-Prot#:Q92556
Uniprot	Q92556
GeneID	9844;
Calculated MW	84 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


Application Details

WB: 1:500-1:2,000 IHC: 1:50-1:200 IP: 1:50-1:100FC: 1:50-1:100


Images




Western blot analysis of ELMO1 on mouse spleen tissue and Jurkat cell lysates using anti-ELMO1 antibody at 1/1,000 dilution.



Immunohistochemical analysis of paraffin-embedded rat spleen tissue using anti-ELMO1 antibody. Counter stained with hematoxylin. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins.



Immunohistochemical analysis of paraffin-embedded mouse spleen tissue using anti-ELMO1 antibody. Counter stained with hematoxylin. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins.



Flow cytometric analysis of SH-SY-5Y cells with ELMO1 antibody at 1/50 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

Background

Elmo (engulfment and cell motility) proteins share similarity to *C. elegans* CED-12. The *C. elegans* genes *ced-2*, *ced-5*, *ced-10* and *ced-12*, and their mammalian homologs, CRKII, DOCK1, RAC1 and ELMO, mediate cytoskeletal rearrangements during phagocytosis of apoptotic cells as well as cell motility. Elmo1 associates with DOCK 180 and may influence phagocytosis and effect cell shape changes. Src family kinase-mediated tyrosine phosphorylation of Elmo1 influences signaling through Elmo1/Crk/DOCK 180 pathways. Elmo2 interacts directly with Rho G in a GTP-dependent manner and forms a ternary complex with DOCK 180 to induce activation of Rac 1. The Rho G-Elmo2-DOCK 180 pathway is required for activation of Rac 1 and cell spreading mediated by integrin, as well as for neurite outgrowth induced by nerve growth factor. Elmo3 acts in association with DOCK 180 and Crk II and may be required in complex with DOCK 180 to activate Rac/Rho small GTPases.

References

Note: This product is for in vitro research use only