Nuf2 Rabbit mAb

Catalog No: #49882

Package Size: #49882-1 50ul #49882-2 100ul



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com


_		4.5	
Des	cri	ntic	าท
レしる	UH	ρ u	ווע

Product Name	Nuf2 Rabbit mAb		
Host Species	Recombinant Rabbit		
Clonality	Monoclonal antibody		
Clone No.	JG34-64		
Purification	ProA affinity purified		
Applications	WB,FC		
Species Reactivity	Hu		
Immunogen Description	Recombinant protein within human Nuf2 aa 1-200.		
Other Names	Cancer/testis antigen 106 antibody CDCA 1 antibody CDCA1 antibody Cell division cycle associated 1 antibody Cell division cycle associated protein 1 antibody Cell division cycle-associated protein 1 antibody CT106 antibody hNuf 2 antibody NUF 2 antibody NUF 2 NDC80 kinetochore complex component antibody NUF 2 NDC80 kinetochore complex component homolog antibody NUF 2, S. cerevisiae, homolog of antibody NUF 2_HUMAN antibody NUF 2 antibody		
Accession No.	Swiss-Prot#:Q9BZD4		
Uniprot	Q9BZD4		
GeneID	83540;		
Calculated MW	54 kDa		
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.		
Storage	Store at -20°C		


Application Details

WB: 1:500-1:2,000 FC: 1:50-1:100

Images



Western blot analysis of Nuf2 on different cell lysates using anti-Nuf2 at 1/1,000 dilution. Positive controlo Ω ½o Ω ½ Line 1: K562 Line 2: PC-3M Line 3: THP-1



Flow cytometric analysis of A431 cells with Nuf2 antibody at 1/50 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

Background

This gene encodes a protein that is highly similar to yeast Nuf2, a component of a conserved protein complex associated with the centromere. Yeast Nuf2 disappears from the centromere during meiotic prophase when centromeres lose their connection to the spindle pole body, and plays a regulatory role in chromosome segregation. The encoded protein is found to be associated with centromeres of mitotic HeLa cells, which suggests that this protein is a functional homolog of yeast Nuf2. Acts as a component of the essential kinetochore-associated NDC80 complex, which is required for chromosome segregation and spindle checkpoint activity. Required for kinetochore integrity and the organization of stable microtubule binding sites in the outer plate of the kinetochore. The NDC80 complex synergistically enhances the affinity of the SKA1 complex for microtubules and may allow the NDC80 complex to track depolymerizing microtubules.

References

Note: This product is for in vitro research use only