

## USP9x Rabbit mAb

Catalog No: #49883



Package Size: #49883-1 50ul #49883-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	USP9x Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JG35-11
Purification	ProA affinity purified
Applications	WB,ICC,IF,IHC,FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Recombinant protein within human USP9X aa 1-200.
Other Names	Deubiquitinating enzyme FAF X antibody Deubiquitinating enzyme FAF-X antibody DFFRX antibody Drosophila fat facets related X linked antibody FAF antibody Fafl antibody Fam antibody Fat facets homolog antibody Fat facets in mammals antibody Fat facets protein related X linked antibody Fat facets protein related, X-linked antibody Fat facets protein-related antibody hFAM antibody MRX99 antibody Probable ubiquitin carboxyl terminal hydrolase FAF X antibody Probable ubiquitin carboxyl-terminal hydrolase FAF-X antibody Ubiquitin carboxyl-terminal hydrolase FAM antibody Ubiquitin specific peptidase 9 X linked antibody Ubiquitin specific peptidase 9, X-linked antibody Ubiquitin specific processing protease FAF X antibody Ubiquitin specific protease 9 X chromosome antibody Ubiquitin thioesterase FAF X antibody Ubiquitin thiolesterase FAF X antibody Ubiquitin thiolesterase FAF-X antibody Ubiquitin-specific protease 9 antibody Ubiquitin-specific-processing protease FAF-X antibody USP9 (gene name) antibody Usp9x antibody USP9X_HUMAN antibody Uubiquitin specific protease 9, X chromosome (fat facets like Drosophila) antibody X chromosome antibody X-linked antibody
Accession No.	Swiss-Prot#:Q93008
Uniprot	Q93008
GeneID	8239;
Calculated MW	292 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:1,000-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200FC: 1:50-1:100


## Images




Western blot analysis of USP9X on different lysates using anti-USP9X antibody at 1/1,000 dilution. Positive control: Lane 1: SiHa Lane 2: A549 Lane 3: 293 Lane 4: Mouse colon




Immunohistochemical analysis of paraffin-embedded human tonsil tissue using anti-USP9X antibody. Counter stained with hematoxylin. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins.




Immunohistochemical analysis of paraffin-embedded mouse fallopian tube tissue using anti-USP9X antibody. Counter stained with hematoxylin. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH6) for 20 mins.




ICC staining USP9X in 293T cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining USP9X in F9 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining USP9X in SiHa cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of 293T cells with USP9X antibody at 1/100 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

## Background

Deubiquitinase involved both in the processing of ubiquitin precursors and of ubiquitinated proteins. May therefore play an important regulatory role at the level of protein turnover by preventing degradation of proteins through the removal of conjugated ubiquitin. Specifically hydrolyzes 'Lys-48', 'Lys-29'- and 'Lys-33'-linked polyubiquitins chains. Essential component of TGF-beta/BMP signaling cascade. Specifically deubiquitinates monoubiquitinated SMAD4, opposing the activity of E3 ubiquitin-protein ligase TRIM33. Deubiquitinates alkylation repair enzyme ALKBH3. OTUD4 recruits USP7 and USP9X to stabilize ALKBH3, thereby promoting the repair of alkylated DNA lesions. Regulates chromosome alignment and segregation in mitosis by regulating the localization of BIRC5/survivin to mitotic centromeres. Involved in axonal growth and neuronal cell migration.

## References

Note: This product is for in vitro research use only