

## CYP27A1 Rabbit mAb

Catalog No: #49968



Package Size: #49968-1 50ul #49968-2 100ul

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com


## Description

Product Name	CYP27A1 Rabbit mAb
Host Species	Recombinant Rabbit
Clonality	Monoclonal antibody
Clone No.	JG40-36
Purification	ProA affinity purified
Applications	WB,ICC,IHC,FC
Species Reactivity	Hu, Ms, Rt
Immunogen Description	Recombinant protein corresponding to human CYP27A1 300-500aa.
Other Names	12-alpha-triol 27-hydroxylase antibody 5-beta-cholestane-3-alpha antibody 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha-triol 26-hydroxylase antibody 5-beta-cholestane-3-alpha, 7-alpha, 12-alpha-triol 27-hydroxylase antibody 5-beta-cholestane-3-alpha,7-alpha,12-alpha-triol 27-hydroxylase antibody 7-alpha antibody Cholestanetriol 26 monooxygenase antibody CP27 antibody CP27A_HUMAN antibody CTX antibody CYP antibody CYP27 antibody CYP27A1 antibody Cytochrome P 450C27/25 antibody Cytochrome P-450C27/25 antibody Cytochrome P450 27 antibody Cytochrome P450 27 mitochondrial antibody Cytochrome P450 family 27 subfamily A member 1 antibody Cytochrome P450 family 27 subfamily A polypeptide 1 antibody Cytochrome P450 subfamily XXVIIA (steroid 27-hydroxylase cerebrotendinous xanthomatosis) polypeptide 1 antibody mitochondrial antibody Sterol 26 hydroxylase antibody Sterol 26 hydroxylase mitochondrial antibody Sterol 26-hydroxylase antibody Sterol 27 hydroxylase antibody Sterol 27-hydroxylase antibody Vitamin D(3) 25 hydroxylase antibody Vitamin D(3) 25-hydroxylase antibody
Accession No.	Swiss-Prot#:Q02318
Uniprot	Q02318
GeneID	1593;
Calculated MW	60 kDa
Formulation	1*TBS (pH7.4), 1%BSA, 40%Glycerol. Preservative: 0.05% Sodium Azide.
Storage	Store at -20°C


## Application Details

WB: 1:500-1:2,000 IHC: 1:50-1:200 ICC: 1:50-1:200 FC: 1:50-1:100


## Images




Western blot analysis of CYP27A1 on different tissue lysates using anti-CYP27A1 antibody at 1/500 dilution. Positive control: Lane 1: Rat heart Lane 2: Human liver Lane 2: Human kidney




Immunohistochemical analysis of paraffin-embedded human liver cancer tissue using anti-CYP27A1 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human kidney tissue using anti-CYP27A1 antibody. Counter stained with hematoxylin.




Immunohistochemical analysis of paraffin-embedded human small intestine tissue using anti-CYP27A1 antibody. Counter stained with hematoxylin.



ICC staining CYP27A1 in HepG2 cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



ICC staining CYP27A1 in LOVO cells (green). The nuclear counter stain is DAPI (blue). Cells were fixed in paraformaldehyde, permeabilised with 0.25% Triton X100/PBS.



Flow cytometric analysis of HepG2 cells with CYP27A1 antibody at 1/100 dilution (purple) compared with an unlabelled control (cells without incubation with primary antibody; yellow). Alexa Fluor 488-conjugated goat anti-rabbit IgG was used as the secondary antibody.

## Background

P450 enzymes constitute a family of monooxygenase enzymes that are involved in the metabolism of a wide array of endo-genous and xenobiotic compounds. P450 enzymes can be classified, based on their sequence similarities, into distinct subfamilies, which include CYP1A and CYP2A. Other P450 family members include CYP19, also designated aromatase (P450arom), which catalyzes the conversion of C19 steroids to estrogens in various tissues, including placenta, gonads, adipose tissue, skin and brain. CYP19 expression is controlled by hormonally regulated promoters in different tissues and increased aromatase activity is associated with familial gynecomastia. Also, a polymorphic allele of CYP19 (repeat (TTTA)<sub>12</sub>) is present in a majority of breast cancer patients. P450 cholesterol 7 $\alpha$ -hydroxylase, CYP7A1, is the rate limiting enzyme of bile acid synthesis in the liver, and its expression is mediated by the bile acid receptor FXR. CYP27A1 catalyzes vitamin D3 25-hydroxylation and is localized to the mitochondria in kidney and liver.

## References

Note: This product is for in vitro research use only