

# Bovine Troponin T, fast skeletal muscle (TNNT3) ELISA Kit



Catalog No: #EK6136

Orders: order@signalwayantibody.com

Package Size: #EK6136-1 48T #EK6136-2 96T

Support: tech@signalwayantibody.com

## Description

Product Name	Bovine Troponin T, fast skeletal muscle (TNNT3) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Bovine (Bos taurus; Cattle)
Other Names	DKFZp779M2348; TNTF; OTTHUMP00000014435 troponin T; fast skeletal muscle troponin T3; skeletal; fast troponin-T3; skeletal; fast
Accession No.	Q8MKI3
Uniprot	Q8MKI3
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.  The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

## Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

## Product Description

**Detection Method:**SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TNNT3 in samples. An antibody specific for TNNT3 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTNNT3 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TNNT3 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TNNT3 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**Subunit T binds the troponin complex to the tropomyosin complex and is also required for Ca(2+)-mediated activation of actomyosin ATPase activity. There are 3 different troponin T genes that encode tissue-specific isoforms of subunit T for fast skeletal-, slow skeletal-, and cardiac-muscle. This gene encodes fast skeletal troponin T protein; also known as troponin T type 3. Alternative splicing results in multiple transcript variants encoding additional distinct troponin T type 3 isoforms. A developmentally regulated switch between fetal/neonatal and adult troponin T type 3 isoforms occurs. Additional splice variants have been described but their biological validity has not been established. Mutations in this gene may cause distal arthrogyrosis multiplex congenita type 2B (DA2B).

---

Note: This product is for in vitro research use only