

Mouse Cluster of differentiation 30 (CD30) ELISA Kit

Catalog No: #EK6214



Package Size: #EK6214-1 48T #EK6214-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Mouse Cluster of differentiation 30 (CD30) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	CD30; D1S166E; Ki-1; CD30 antigen CD30L receptor Ki-1 antigen cytokine receptor CD30 lymphocyte activation antigen CD30
Accession No.	Q60846
Uniprot	Q60846
GeneID	21941;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:15.6-1000 pg/mL

Sensitivity:3.9 pg/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:Sandwich**Test principle:**This assay employs a two-site sandwich ELISA to quantitate TNFRSF8 in samples. An antibody specific for TNFRSF8 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTNFRSF8 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TNFRSF8 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TNFRSF8 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:** This protein is a member of the TNF-receptor superfamily. This receptor is expressed by activated, but not by resting, T and B cells. TRAF2 and TRAF5 can interact with this receptor, and mediate the signal transduction that leads to the activation of NF-kappaB. This receptor is a positive regulator of apoptosis, and also has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. Two alternatively spliced transcript variants of this gene encoding distinct isoforms have been reported.

Note: This product is for in vitro research use only