Human Transmembrane protein 9B (TMEM9B) ELISA Kit

Catalog No: #EK6345

Package Size: #EK6345-1 48T #EK6345-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Human Transmembrane protein 9B (TMEM9B) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	C11orf15;
Accession No.	Q9NQ34
Uniprot	Q9NQ34
GeneID	56674;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details Detect Range:Request Information Sensitivity:Request Information Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TMEM9B in samples. An antibody specific for TMEM9B has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTMEM9B present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TMEM9B is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TMEM9B bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:The intracellular signaling pathway by which TNF induces its pleiotropic actions is well-characterized and includes unique components, as well as modules shared with other signaling pathways. TMEM9B as an important component of TNF signaling and which is a module shared with the IL-1β and Toll-like receptor (TLR) pathways. is a glycosylated protein localized in membranes of the lysosome and partially in early endosomes. The expression of TMEM9B is required for the production of proinflammatory cytokines induced by TNF, IL-1β, and TLR ligands, but not for apoptotic cell death triggered by TNF or Fas ligand. TMEM9B is essential in TNF activation of both the NF-κB and MAPK pathways.TMEM9B is a key component of inflammatory signaling pathways and suggest that endosomal or lysosomal compartments regulate these pathways.

Note: This product is for in vitro research use only