Human Transmembrane protein 168 (TMEM168) ELISA Kit

Catalog No: #EK6426

Package Size: #EK6426-1 48T #EK6426-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Human Transmembrane protein 168 (TMEM168) ELISA Kit
ELISA Kit
ELISA
Human (Homo sapiens)
DKFZp564C012; FLJ13576; OTTHUMP00000207378
Q9H0V1
Q9H0V1
64418;
The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details Detect Range:Request Information Sensitivity:Request Information Sample Type:Serum, Plasma, Other biological fluids Sample Volume: 1-200 µL Assay Time:1-4.5h Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TMEM168 in samples. An antibody specific for TMEM168 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTMEM168 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TMEM168 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TMEM168 bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:TMEM168 is a transmembrane protein (TP) protein that goes from one side of a membrane through to the other side of the membrane. Many TPs function as gateways or "loading docks" to deny or permit the transport of specific substances across the biological membrane, to get into the cell, or out of the cell as in the case of waste byproducts. As a response to the shape of certain molecules these "freight handling" TPs may have special ways of folding up or bending that will move a substance through the biological membrane. A transmembrane protein is a polytopic protein that spans an entire biological membrane. Transmembrane proteins aggregate and precipitate in water. They require detergents or nonpolar solvents for extraction, although some of them (beta-barrels) can be also extracted using denaturing agents.

Note: This product is for in vitro research use only