

Human Tomoregulin-2 (TMEFF2) ELISA Kit

Catalog No: #EK6454



Package Size: #EK6454-1 48T #EK6454-2 96T

Orders: order@signalwayantibody.com

Support: tech@signalwayantibody.com

Description

Product Name	Human Tomoregulin-2 (TMEFF2) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	HPP1; TENB2; TPEF; TR; tomoregulin transmembrane protein TENB2
Accession No.	Q9UIK5
Uniprot	Q9UIK5
GeneID	23671;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TMEFF2 in samples. An antibody specific for TMEFF2 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTMEFF2 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TMEFF2 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TMEFF2 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**The deduced 368-amino acid TRa protein contains 2 follistatin-like domains and a conserved EGF-like domain. It also has a predicted N-terminal signal peptide, a transmembrane domain, a cytosolic tail G protein-activating motif, and potential sites for N-linked glycosylation and glycosaminoglycan attachment. Unlike other EGF/NRG family members, TRa has a histidine replacing a conserved arginine essential for EGF receptor recognition. The TRb and TRc proteins contain 418 and 379 amino acids, respectively, and differ from one another and TRa in the cytoplasmic domain. Northern blot analysis detected expression of approximately 2.2- and 3.1-kb TR transcripts predominantly in brain. The 374-amino acid TMEFF2 protein shares approximately 36% and 99% sequence identity with TMEFF1 and mouse Tmeff2, respectively.

Note: This product is for in vitro research use only