## Mouse Toll-like receptor 8 (TLR8) ELISA Kit

Catalog No: #EK6495

Package Size: #EK6495-1 48T #EK6495-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Descri	nti	Or	า
	IJι	U	ш

Product Name	Mouse Toll-like receptor 8 (TLR8) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	CD288; MGC119599; MGC119600;
Accession No.	P58682
Uniprot	P58682
GeneID	170744;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage
	at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

## **Application Details**

Detect Range:0.312-20 ng/mL	
Sensitivity:0.108 ng/mL	
Sample Type:Serum, Plasma, Other biological fluids	
Sample Volume: 1-200 μL	
Assay Time:1-4.5h	
Detection wavelength:450 nm	

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TLR8 in samples. An antibody specific for TLR8 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTLR8 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TLR8 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TLR8 bound in the initial step. The color development is stopped and the intensity of the color is measured. Product Overview: Toll-like receptor 8 is a member of the Toll-like receptor (TLR) family which plays a fundamental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Drosophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytokines necessary for the development of effective immunity.

The various TLRs exhibit different patterns of expression. This gene is predominantly expressed in lung and peripheral blood leukocytes, and lies in close proximity to another family member, TLR7, on chromosome X.TLR8 recognises G-rich oligonucleotides.

Note: This product is for in vitro research use only