

Human Protein phosphatase Slingshot homolog 1 (SSH1) ELISA Kit



Catalog No: #EK6663

Orders: order@signalwayantibody.com

Package Size: #EK6663-1 48T #EK6663-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Human Protein phosphatase Slingshot homolog 1 (SSH1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Human (Homo sapiens)
Other Names	FLJ21928; FLJ38102; KIAA1298; SSH1L; slingshot 1
Accession No.	Q8WYL5
Uniprot	Q8WYL5
GeneID	54434;
Storage	<p>The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition.</p> <p>The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).</p>

Application Details

Detect Range:0.312-20 ng/mL

Sensitivity:0.105 ng/mL

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 µL

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate SSH1 in samples. An antibody specific for SSH1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anySSH1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for SSH1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of SSH1 bound in the initial step. The color development is stopped and the intensity of the color is measured.**Product Overview:**The ADF (actin-depolymerizing factor)/cofilin family is composed of stimulus-responsive mediators of actin dynamics. ADF/cofilin proteins are inactivated by kinases such as LIM domain kinase-1 (LIMK1). The SSH family appears to play a role in actin dynamics by reactivating ADF/cofilin proteins in vivo. Transcript analysis suggested that at least 6 polypeptides would be produced from these genes. SSH1 encodes 3 isoforms, SSH1L (1,049 amino acids), SSH1S (692 amino acids), and SSH1B (148 amino acids), and SSH2 encodes 2 isoforms, SSH2 (449 amino acids) and SSH2B (195 amino acids). The SSH3 protein has 471 amino acids. Besides the catalytic domain, 2 other domains are conserved between Drosophila ssh and the human SSHs and are unique to the SSH family.

Note: This product is for in vitro research use only