WT1 antibody

Catalog No: #22694




Orders: order@signalwayantibody.com


Description	Support: tech@signalwayantibody.com
Product Name	WT1 antibody
Host Species	Rabbit
Clonality	Polyclonal
Purification	Purified by antigen-affinity chromatography.
Applications	WB IHC IF
Species Reactivity	Hu
Immunogen Type	Recombinant Protein
Immunogen Description	Recombinant protein fragment contain a sequence corresponding to a region within amino acids 277 and 495
	(P19544) of Wilms Tumor 1
Target Name	WT1
Accession No.	Swiss-Prot:P19544Gene ID:7490
Uniprot	P19544
GenelD	7490;
Concentration	1.09mg/ml
Formulation	Supplied in 0.1M Tris-buffered saline with 10% Glycerol (pH7.0). 0.01% Thimerosal was added as a
	preservative.
Storage	Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details Predicted MW: 56kd Western blotting: 1:500-1:3000 Immunohistochemistry: 1:50-1:500 Immunofluorescence: 1:100-1:200


Images



Sample (30 ug of whole cell lysate) A: Molt-4 B: Raji 10% SDS PAGE Primary antibody diluted at 1: 1000



Immunohistochemical analysis of paraffin-embedded RT2 xenograft, using Wilms Tumor 1 antibody at 1: 500 dilution.



Immunofluorescence analysis of paraformaldehyde-fixed HeLa, using WT1 antibody at 1: 200 dilution.

Background

This gene encodes a transcription factor that contains four zinc-finger motifs at the C-terminus and a proline/glutamine-rich DNA-binding domain at the N-terminus. It has an essential role in the normal development of the urogenital system, and it is mutated in a small subset of patients with Wilm's tumors. Multiple transcript variants, resulting from alternative splicing at two coding exons, have been well characterized. There is also evidence for the use of non-AUG (CUG) translation initiation site upstream of, and in-frame with the first AUG, leading to additional isoforms. Authors of PMID:7926762 also provide evidence that WT1 mRNA undergoes RNA editing in human and rat, and that this process is tissue-restricted and developmentally regulated. [provided by RefSeq]

Note: This product is for in vitro research use only