ACTR-II antibody

Catalog No: #22695

Description Product Name

Host Species Clonality



Orders: order@signalwayantibody.com Support: tech@signalwavantibodv.com

	Support. tech@signaiwayantibody.com
ACTR-II antibody	
Rabbit	
Polyclonal	
Purified by antigen-affinity chromatography.	


Purified by antigen-affinity chromatography.
WB
Hu
Recombinant protein
Recombinant protein fragment contain a sequence corresponding to a region within amino acids 267 and 509
of Human ACVR2A
ACTR-II
Swiss-Prot:P27037Gene ID:92
P27037
92;
0.9mg/ml
Supplied in 0.1M Tris-buffered saline with 20% Glycerol (pH7.0). 0.01% Thimerosal was added as a
preservative.
Store at -20°C for long term preservation (recommended). Store at 4°C for short term use.

Application Details

Predicted MW: 58kd

Western blotting: 1:500-1:3000

Images



Sample (30 ug of whole cell lysate) A: Hela B: Hep G2 7.5% SDS PAGE Primary antibody diluted at 1: 10000

Background

This gene encodes activin A type II receptor. Activins are dimeric growth and differentiation factors which belong to the transforming growth factor-beta (TGF-beta) superfamily of structurally related signaling proteins. Activins signal through a heteromeric complex of receptor serine kinases which

include at least two type I (I and IB) and two type II (II and IIB) receptors. These receptors are all transmembrane proteins, composed of a ligand-binding extracellular domain with cysteine-rich region, a transmembrane domain, and a cytoplasmic domain with predicted serine/threonine specificity. Type I receptors are essential for signaling; and type II receptors are required for binding ligands and for expression of type I receptors. Type I and II receptors form a stable complex after ligand binding, resulting in phosphorylation of type I receptors by type II receptors. Type II receptors are considered to be constitutively active kinases. [provided by RefSeq]

Note: This product is for in vitro research use only