

Pig Transforming growth factor-beta-induced protein ig-h3 (TGFBI) ELISA Kit



Catalog No: #EK7074

Orders: order@signalwayantibody.com

Package Size: #EK7074-1 48T #EK7074-2 96T

Support: tech@signalwayantibody.com

Description

Product Name	Pig Transforming growth factor-beta-induced protein ig-h3 (TGFBI) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Pig (<i>Sus scrofa</i> ; Porcine)
Other Names	BIGH3; CDB1; CDG2; CDGG1; CSD; CSD1; CSD2; CSD3; EBMD; LCD1; RGD-containing collagen-associated protein kerato-epithelin
Accession No.	O11780
Uniprot	O11780
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5% within the expiration date under appropriate storage condition. The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days, and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

Application Details

Detect Range:Request Information

Sensitivity:Request Information

Sample Type:Serum, Plasma, Other biological fluids

Sample Volume: 1-200 μ L

Assay Time:1-4.5h

Detection wavelength:450 nm

Product Description

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate TGFBI in samples. An antibody specific for TGFBI has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anyTGFBI present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for TGFBI is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of TGFBI bound in the initial step. The color development is stopped and the intensity of the color is measured.Product Overview:TGF- β is capable of producing a variety of effects and virtually all cell types respond to this factor in some way. The inappropriate presence of active TGF- β 1 has been implicated in a variety of pathological conditions Because of the necessity for regulating its activity tightly, TGF- β is secreted by cells in the form of an inactive complex. This complex consists of TGF- β 1 associated non-covalently with a protein designated the latency associated peptide (LAP).

TGF- β 1 and LAP represent components of a pro-peptide that is cleaved in a post-golgi compartment prior to secretion. LAP and TGF- β 1 each consist of a disulfide-linked homodimer and the association of these two components renders TGF- β 1 inactive and inaccessible to anti-TGF- β antibodies.

Note: This product is for in vitro research use only