## Mouse Son of sevenless homolog 1 (SOS1) ELISA Kit

Catalog No: #EK7194

Package Size: #EK7194-1 48T #EK7194-2 96T



Orders: order@signalwayantibody.com Support: tech@signalwayantibody.com

Description	
Product Name	Mouse Son of sevenless homolog 1 (SOS1) ELISA Kit
Brief Description	ELISA Kit
Applications	ELISA
Species Reactivity	Mouse (Mus musculus)
Other Names	GF1; GGF1; GINGF; HGF; NS4; OTTHUMP00000128306 gingival fibromatosis; hereditary; 1 guanine
	nucleotide exchange factor son of sevenless homolog 1
Accession No.	Q62245
Uniprot	Q62245
GeneID	20662;
Storage	The stability of ELISA kit is determined by the loss rate of activity. The loss rate of this kit is less than 5%
	within the expiration date under appropriate storage condition.
	The loss rate was determined by accelerated thermal degradation test. Keep the kit at 37C for 4 and 7 days,
	and compare O.D.values of the kit kept at 37C with that of at recommended temperature. (referring from China
	Biological Products Standard, which was calculated by the Arrhenius equation. For ELISA kit, 4 days storage

## **Application Details**

Detect Range:Request Information		
Sensitivity:Request Information		
Sample Type:Serum, Plasma, Other biological fluids		
Sample Volume: 1-200 μL		
Assay Time:1-4.5h		
Detection wavelength:450 nm		

at 37C can be considered as 6 months at 2 - 8C, which means 7 days at 37C equaling 12 months at 2 - 8C).

## **Product Description**

Detection Method:SandwichTest principle:This assay employs a two-site sandwich ELISA to quantitate SOS1 in samples. An antibody specific for SOS1 has been pre-coated onto a microplate. Standards and samples are pipetted into the wells and anySOS1 present is bound by the immobilized antibody. After removing any unbound substances, a biotin-conjugated antibody specific for SOS1 is added to the wells. After washing, Streptavidin conjugated Horseradish Peroxidase (HRP) is added to the wells. Following a wash to remove any unbound avidin-enzyme reagent, a substrate solution is added to the wells and color develops in proportion to the amount of SOS1 bound in the initial step. The color development is stopped and the intensity of the color is measured.

Note: This product is for in vitro research use only